

The Role of the Endocannabinoid System in Psychiatric Disorders and Symptoms: a Pharmacological fMRI study.

No registrations found.

Ethical review	Positive opinion
Status	Recruiting
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON20111

Source

Nationaal Trial Register

Brief title

THC-phMRI in patients

Health condition

psychiatric disorders (schizophrenia, addiction, OCD, ADHD, depression), fMRI, THC, cannabinoid system

Sponsors and support

Primary sponsor: Prof. Dr. N.F. Ramsey

Source(s) of monetary or material Support: Top Institue Pharma (TI Pharma)

Intervention

Outcome measures

Primary outcome

The main study parameter is the blood oxygen level dependent (BOLD) signal.

Secondary outcome

Behavioral parameters (two sets of visual analogue scales), cerebral blood flow (measured with Arterial Spin Labeling), the concentration of plasma THC and its main metabolites, and the performance on neuropsychological tests will be measured.

Study description

Background summary

Extensive research has identified and characterized an endocannabinoid system in the human central nervous system. It consists of cannabinoid receptors and endocannabinoids that work on the receptors. Recently, the endocannabinoid system has been associated with several psychiatric disorders, such as schizophrenia, depression and addiction. This system provides a novel potential target for medical treatment of these disorders. This study assesses the role of the endocannabinoid system in symptoms of psychiatric disorders. THC, which is the main psychoactive constituent of cannabis, works on cannabinoid receptors as an agonist. The study is a randomized, double-blind, placebo-controlled pharmacological MRI experiment in which healthy controls will be compared to patients with a psychiatric disorder. The study consists of two test days, on which subjects receive either placebo or THC by means of a vaporizer. On a test day subjects will perform three cognitive tasks, during which functional MRI scans will be measured. Also, a series of neuropsychological tests will be performed. Before every task subjects receive a new dose of THC or placebo.

Study objective

The endocannabinoid system is involved in cognitive functioning, and therefore, challenging this system with THC will influence cognitive processing in humans. Moreover, subjects with cognitive impairments (i.e. psychiatric patients) will react differently on a cannabinoid challenge than healthy controls.

Study design

Functional MRI scans will be obtained between 7 and 21 minutes after inhalation. Behavioral parameters will be measured and blood samples will be withdrawn before and after functional MRI. Two testdays are performed (one day placebo, one day THC), two weeks apart.

Intervention

1. THC will be inhaled (5 times: first dose 6 mg, upload doses 1 mg) by means of a Vulcano Vaporizer;

2. Functional MRI scans will be performed during cognitive tasks.

Contacts

Public

H.H. Hell, van
University Medical Center Utrecht,
Department of Neurology and Neurosurgery,
Heidelberglaan 100
Utrecht 3584 CX
The Netherlands
+31 (0)88 7555873

Scientific

H.H. Hell, van
University Medical Center Utrecht,
Department of Neurology and Neurosurgery,
Heidelberglaan 100
Utrecht 3584 CX
The Netherlands
+31 (0)88 7555873

Eligibility criteria

Inclusion criteria

1. Healthy control or Diagnosed with one psychiatric disorder, according to DSM-IV criteria, axis I:
 - A. Schizophrenia;
 - B. Depression;
 - C. ADHD;
 - D. Addiction to nicotine (heavy smokers);
 - E. OCD.
2. Male;
3. Current occasional cannabis use since at least one year (<1/week and \geq 4/year) without

known negative implications (e.g. bad trip, cannabis-induced psychosis);

4. Right-handedness, assessed with the Edinburgh Handedness Inventory (Oldfield, 1971);

5. Written informed consent of the subject.

Exclusion criteria

1. Clinical significant abnormalities, except for the predetermined psychiatric disorder;

2. For healthy controls, first degree relatives with a psychiatric disorder according to DSM IV;

3. Impaired physical health evaluated by medical history and physical (including neurological) examination;

4. Current diagnosis of abuse of drugs or alcohol (according to DSM-IV) except for tobacco;

5. Past but recent diagnosis of abuse of drugs or alcohol other than tobacco, i.e. within 12 months preceding study inclusion;

6. Body Mass Index (B.M.I.) $<18 \text{ kg/m}^2$ or $>28 \text{ kg/m}^2$;

7. Any subject who received any investigational medication within 90 days prior to the start of the study or who is scheduled to receive an investigational drug;

8. The use of any medication within three weeks prior to the start of the study, except for paracetamol and medication for the psychiatric disorder;

9. Blood donation within 3 months before the start of the study;

10. Claustrophobia;

11. Metal objects in or around the body (braces, pacemaker, metal fragments).

Study design

Design

Study type:	Interventional
Intervention model:	Crossover
Allocation:	Randomized controlled trial

Masking:	Double blinded (masking used)
Control:	Placebo

Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-02-2009
Enrollment:	120
Type:	Anticipated

Ethics review

Positive opinion	
Date:	29-04-2009
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL1685
NTR-old	NTR1787
Other	METC Utrecht : 07/371
ISRCTN	ISRCTN wordt niet meer aangevraagd

Study results

Summary results

N/A