

# Time to repeat SLT in glaucoma patients

No registrations found.

<b>Ethical review</b>	Not applicable
<b>Status</b>	Recruiting
<b>Health condition type</b>	-
<b>Study type</b>	Observational non invasive

## Summary

### ID

NL-OMON20154

### Source

Nationaal Trial Register

### Brief title

SLT

### Health condition

Glaucoma, laser treatment, glaucoom, laser behandeling

## Sponsors and support

**Primary sponsor:** none

**Source(s) of monetary or material Support:** none

## Intervention

## Outcome measures

### Primary outcome

time to redo SLT

### Secondary outcome

IOP control after primary and secondary SLT

# Study description

## Background summary

SLT has proven to be a viable alternative for medication in the treatment of glaucoma patients. Like argon laser trabeculoplasty, its effect seems to lower in time. Its repeatability is therefore important. We aim to investigate how long it takes for the effect of SLT wears off and whether a second SLT is as effective as the first SLT to lower IOP.

## Study objective

- 1, the effect of SLT diminishes in time; how fast does IOP get below target IOP after SLT?
- 2, is a repeat SLT effective to get/keep IOP under target IOP?
- 3, is there a racial difference in effect of SLT?

## Study design

baseline, 1 week, 3, 6, 12, 18, 24, 30, 36, 42, 48, 60 months

## Intervention

- 1, redo of SLT when target IOP is exceeded by more than 2 mmHg when IOP was controlled at least 6 months after initial SLT
- 2, control group under medication

# Contacts

## Public

Den Brem 93

Myrjam de Keyser  
Wilrijk 2610  
Belgium  
00 32 473/89 30 01

## Scientific

Den Brem 93

Myrjam de Keyser  
Wilrijk 2610  
Belgium

## Eligibility criteria

### Inclusion criteria

Patients with ocular hypertension or open angle glaucoma. Other types of glaucoma than open angle glaucoma, corneal disease that inhibit good visualization of the trabecular meshwork, taking of systemic or local steroids.

### Exclusion criteria

Other types of glaucoma than open angle glaucoma, corneal disease that inhibit good visualization of the trabecular meshwork, taking of systemic or local steroids.

## Study design

### Design

Study type:	Observational non invasive
Intervention model:	Parallel
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)
Control:	Active

### Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-01-2017
Enrollment:	200
Type:	Anticipated

## Ethics review

Not applicable

Application type:

Not applicable

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL6144
NTR-old	NTR6299
Other	: NTR5417

## Study results

### Summary results

De Keyser M, De Belder M, De Belder S and De Groot V. Where does selective laser trabeculoplasty stand now? A review. Eye and Vision. 2016;3:10

DOI 10.1186/s40662-016-0041-y

De Keyser M. Power is not energy. Letter to the Editor. Journal of Glaucoma, May 5, 2016.

Doi:10.1097/IJG.0000000000000432

De Keyser M, De Belder M, De Groot V. Randomized prospective study of the use of anti-inflammatory drops after selective laser trabeculoplasty. Journal of Glaucoma, 26(2), e22-e29.doi:10.1097/IJG.0000000000000522.

De Keyser M, De Belder M, De Groot V. Selective laser trabeculoplasty in pseudophakic and phakic eyes: a prospective study. International Journal of Ophthalmology, 10;4:2017.

De Keyser M, De Belder M, De Groot V. Prospective study on the effect of selective laser trabeculoplasty in normal tension glaucoma. International Journal of Ophthalmology and Eye Science, S1:008, 36-41, 2016. IJOES-2332-290X-S1-008

De Keyser M, De Belder M, De Groot V. Treatment-related quality of life in glaucoma patients: comparison between selective laser trabeculoplasty and medication. International Journal of Ophthalmology, 10; 4: 2017.

De Keyser M, De Belder M, De Belder J, De Groot V. Effect of selective laser trabeculoplasty in glaucoma patients with high or low central corneal thickness. Insights in Ophthalmology, vol 1, 1:3, 2017.

De Keyser M, De Belder M, De Belder J, De Groot V. Selective laser trabeculoplasty as replacement therapy in medically controlled glaucoma patients. In submission.