

Bipolar Radiofrequency(RF) Lesion of the thoracic medial branch of the posterior ramus for treatment of posterior mechanical spinal pain

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON21338

Source

Nationaal Trial Register

Brief title

BiRThoFP

Health condition

thoracic posterior mechanical (facet) pain
bipolar RF lesion , thoracic facet pain

Sponsors and support

Primary sponsor: None

Source(s) of monetary or material Support: Neurotherm for support in writing the manuscript

Intervention

Outcome measures

Primary outcome

VAS score

PDI (Pain Disability Index)

Secondary outcome

global perceived effect

willingness for repetition of treatment

Study description

Background summary

In the general population the prevalence of thoracic pain is estimated to be 13%¹. In a population with localized thoracic pain the prevalence of thoracic facet joint pain is found to be 42%²

When thoracic facet joint pain proves to be refractory to conventional treatment (medication, physical exercise, TENS) interventional management may be considered. In a recently published randomized controlled trial the effect of infiltration of the medial branch of the thoracic dorsal ramus with local anesthetic with or without corticosteroid was investigated. ³ At 2 years follow-up 80% and 84% of the patients in respectively the group with LA alone and the group treated with LA + corticosteroids. These findings clearly indicate the role of the thoracic facet joints in thoracic pain. Closer analysis of the information, however, shows that patients received a mean of 5.9 injections over the two year study period and the mean duration of pain relief was 18.9 weeks.

Radiofrequency (RF) treatment of the medial branch of the dorsal ramus is well documented for the management of lumbar facet joint pain.⁴ There is less evidence on the use of RF of the medial branch of the thoracic ramus dorsalis. Positive results up to 2 months after the intervention are documented.⁵ It has been suggested that increasing the size of the lesion would result in better treatment outcome. ⁶ Therefore, we assume that using bipolar RF treatment will increase the lesion size and thus improve treatment outcome.

The advantage of bipolar RF over other techniques used to increase the lesion size, such as cooled RF is a better delineation of the lesion, that occurs between the two electrodes.

Study objective

Bipolar radiofrequency lesioning of the thoracic medial branch of the dorsal ramus is more effective than a monopolar radiofrequency lesion

Study design

Outcome will be measured 2 months after the procedure. A 6 month follow-up is also scheduled. The first analysis will however be performed at 2 months.

Intervention

The diagnostic process includes identification of the thoracic levels involved in the pain problem. Because the facet joints are innervated from the adjacent cranial and the same level, and in facet pain mostly at least 2 segmental levels are involved, each intervention is performed at least at 3 levels.

The insertion place of the electrode is identified with an AP projection of the C-arm with the patient in a prone position and a metal ruler to find the proper level. A perfect AP fluoroscopic view shows the end plates of the vertebrae to be neatly projected over each other. The transverse processes are identified and the site of insertion of the needles is marked on the skin. An RF needle is inserted parallel to the C-arm projection until contact is made with the bone at the junction between the processus articularis superius and the processus transversus. The needle is then positioned slightly more cranially just abolishing bone contact. With the first needle in place a second needle is inserted using the direction of the first one. The target position is 1 cm parallel of the first needle, also making contact more laterally on the transverse process. At every level to be treated two needles are placed in the similar manner. In a lateral view with the C arm, the tip of the needles should be posterior to the line that connects the posterior aspects of the foramen intervertebrale.

Subsequently, neurostimulation initially takes place with 50 Hz first, then with 2 Hz. The 2 Hz stimulation causes contraction of the paravertebral muscles at intensities below 0.5 to 0.7 V. After local anesthetic is injected, a 180 second bipolar (dual) RF treatment at 80°C is made.

To summarize, the treatment consists of the application of high frequency current at the medial branch of the posterior ramus of at least 3 adjacent thoracic levels. The current is passed through two electrodes covering the nerve. The anatomic position of the medial branch in the thoracic spine is different from the cervical and lumbar spine, it is running more cranial and lateral over the transverse process. By making a bipolar RF lesion in this way the obtained lesion is larger than when only one electrode is used. The size of the lesion is defined by the space between the two electrodes, the size of the electrodes and the active tip and the RF lesion temperature.

This study is a factorial study aimed at evaluating the effect of bipolar RF treatment when studied in a larger group of patients. There is no control group.

Contacts

Public

Maaslandziekenhuis
Dept. of Pain therapy
Dr. H. Vanderhoffplein, 1
O. Rohof
Maaslandziekenhuis
Dept. of Pain therapy
Dr. H. Vanderhoffplein, 1
Sittard 6162 BG
The Netherlands
+31 46 457 58 86

Scientific

Maaslandziekenhuis
Dept. of Pain therapy
Dr. H. Vanderhoffplein, 1
O. Rohof
Maaslandziekenhuis
Dept. of Pain therapy
Dr. H. Vanderhoffplein, 1
Sittard 6162 BG
The Netherlands
+31 46 457 58 86

Eligibility criteria

Inclusion criteria

- 1.Age 18-75 years
- 2.Almost continuous uni-or bilateral chronic pain in a circumscribed thoracic spinal region without neurological signs
- 3.Uni- or bilateral paravertebral thoracic tenderness in the same region
- 4.On RX -spine normal or only minimal changes

5.Conservative therapy no or incomplete result

6.Having given informed consent

Exclusion criteria

1."Red Flags" like tumor, infection, fracture

2.Anticoagulation therapy, which may not temporarily be stopped

Study design

Design

Study type:	Interventional
Intervention model:	Factorial
Allocation:	Non controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-01-2014
Enrollment:	70
Type:	Anticipated

Ethics review

Positive opinion	
Date:	24-11-2013
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL4026
NTR-old	NTR4294
Other	13-N-130 : METC nummer
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A