

Ultrasonography for diagnosis of scaphoid fractures: A diagnostic prospective study.

No registrations found.

Ethical review	Not applicable
Status	Suspended
Health condition type	-
Study type	Observational non invasive

Summary

ID

NL-OMON25194

Source

Nationaal Trial Register

Health condition

Scaphoid fractures; all types

- distal pole (including distal articular surface and tuberosity)
- waist
- proximal pole

Sponsors and support

Primary sponsor: Academic Medical Centre, Amsterdam - Departments of plastic-, reconstructive & handsurgery, traumatology and radiology

Source(s) of monetary or material Support: Academic Medical Centre, Amsterdam

Intervention

Outcome measures

Primary outcome

The main objective of this study is to evaluate the use of ultrasonography in detection of occult scaphoid fractures. Sensitivity, specificity and positive and negative predictive values

for ultrasonography in comparison with a CT-scan.

Secondary outcome

Subanalysis of different fractures types:

1. Proximal pole fractures;
2. Waist fractures;
3. Distal pole fractures.

Study description

Background summary

N/A

Study objective

Ultrasonography can be used for detection of occult scaphoid fractures.

Study design

1. Initial presentation at emergency room;
2. Within 7 - 10 days after presentation at emergency room.

Intervention

Ultrasonography of wrist and subsequently a wrist CT.

Contacts

Public

Academisch Medisch Centrum

Afdeling Plastische-, reconstructieve- en handchirurgie

Meibergdreef 9
D.S.P. Heidsieck
Amsterdam 1105 AZ
The Netherlands

+31 (0)6 43034103

Scientific

Academisch Medisch Centrum

Afdeling Plastische-, reconstructieve- en handchirurgie

Meibergdreef 9

D.S.P. Heidsieck

Amsterdam 1105 AZ

The Netherlands

+31 (0)6 43034103

Eligibility criteria

Inclusion criteria

1. Clinical suspicion of a scaphoid fracture;
2. Normal hand/wrist radiographs;
3. Age > 18 years.

Exclusion criteria

1. Other fractures in the affected extremity;
2. Multi trauma patients;
3. Open fractures of the hand;
4. Pregnancy;
5. Insufficient understanding of written and spoken Dutch language.

Study design

Design

Study type: Observational non invasive

Intervention model: Parallel

Allocation: Non controlled trial

Control: N/A , unknown

Recruitment

NL

Recruitment status: Suspended

Start date (anticipated): 01-03-2013

Enrollment: 200

Type: Anticipated

Ethics review

Not applicable

Application type: Not applicable

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL3608
NTR-old	NTR3767
Other	ABR : 43332
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A