

Botulinum Neurotoxin (BoNT) as new treatment for functional (psychogenic) jerky movement disorders.

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON26050

Source

Nationaal Trial Register

Health condition

Botulinum neurotoxin (BoNT) has emerged as a useful therapy for several movement disorders associated with muscle overactivity such as dystonia and jerky movement disorders. At least 2–9% of patients seen in movement disorder clinics suffer from movement disorders with a psychogenic origin and a substantial part of them has jerks. These psychogenic jerky movement disorders cannot be accounted for by a known neurologic syndrome.

Sponsors and support

Primary sponsor: Academic Medical Center

University of Amsterdam

PO-box: 22660

1100 DD Amsterdam

Tel: 0031 (0)20 5669111

Source(s) of monetary or material Support: Prinses Beatrix Fonds

PO Box 85810

2508 CM Den Haag

The Netherlands
Phone (+31) 070 - 3607 607
Fax (+31) 070 - 364 84 50
e-mail s.valentim@prinsesbeatrixfonds.nl

Intervention

Outcome measures

Primary outcome

To assess whether treatment with BoNT leads to improvement of psychogenic jerks according to an independent movement disorder specialist assessed with the Clinical Global Impression - Improvement scale.

Secondary outcome

The effect of BoNT on:

1. The severity of jerks according to a movement disorder specialist;
2. Improvement and severity of the jerk according to the patient;
3. Nature and severity of overall dyskinesia;
4. Jerk frequency;
5. Whether patients consider treatment with BoNT effective;
6. Disability;
7. Quality of life;
8. Co-existent psychiatric disorders;
9. The occurrence of adverse reactions;
10. Muscle weakness.

Study description

Background summary

Rationale:

Botulinum neurotoxin (BoNT) has emerged as a useful therapy for several movement disorders associated with muscle overactivity. At least 2–9% of patients seen in movement disorder clinics suffer from movement disorders with a psychogenic origin and a substantial part of them has jerks. These psychogenic jerky movement disorders cannot be accounted for by a known neurologic syndrome. Therapy of psychogenic jerks currently focuses on frequently co-occurring psychiatric disease, but results are poor. In this project, we will study the effect of BoNT on movement disorders of psychogenic origin.

Objective:

To evaluate the effect of treatment with BoNT on psychogenic jerks.

Study design:

A monocenter study consisting of two parts: a double-blind randomized placebo controlled intervention study of 16 weeks and an uncontrolled follow-up study of one year to evaluate the long-term effects of BoNT.

Study population:

Patients with at least one invalidating consistent type of psychogenic jerk that is present for 1 year or longer.

Intervention:

During the trial phase of the study, patients will receive two BoNT or placebo injections with an interval of 3 months. The number of muscles injected and the doses to be administered in an individual patient will be determined by an experienced neurophysiologist analogous to treatment of dystonia. Hereafter, for the follow-up study all patients will receive 4 BoNT injections at intervals of 3 months.

Main study parameters/endpoints:

Primary outcome measure: To assess whether treatment with BoNT leads to improvement of psychogenic jerks according to an independent movement disorder specialist assessed with

the Clinical Global Impression - Improvement scale.

Secondary outcome measures are the effect of BoNT on: the severity of jerks according to a movement disorder specialist; improvement and severity of the jerk according to the patient; nature and severity of overall dyskinesia; jerk frequency; whether patients consider treatment with BoNT effective; disability; quality of life; co-existent psychiatric disorders and the occurrence of adverse reactions; muscle weakness

Nature and extent of the burden and risks associated with participation, benefit and group relatedness:

The risks associated with participation in this study are low: BoNT is considered a safe therapy in other movement disorders. The most common side effects of BoNT are local weakness and pain and side effects are reversible. The neuropsychological and psychiatric questionnaires used in our study are considered to be mildly psychologically stressful.

Study objective

Therapy of psychogenic jerks currently focuses on frequently co-occurring psychiatric disease, but results are poor. In this project, we will study the effect of BoNT on movement disorders of psychogenic origin in a mono centred randomised controlled trial. The study consists of two parts: a double-blind randomized placebo controlled intervention study of 16 weeks and an uncontrolled follow-up study of one year to evaluate the long-term effects of BoNT. Hypothesis is that administration of BoNT is an effective treatment for jerky psychogenic movement disorders that significantly compromise patients.

Study design

Baseline (T = 0):

1. Neurological examination, recorded on video;
2. Psychiatric interview, recorded on video;
3. Questionnaires (psychiatry, disability and tic assessment);
4. CGI – severity by examiner (based on video) and patient;
5. Counting of the amount of jerks during 10 minutes of the psychiatric interview on video;
6. AIMS and PMD scale scores based on the recorded video during neurological examination.

T= 16 weeks:

1. Neurological examination, recorded on video;
2. Psychiatric interview, recorded on video;
3. Questionnaires (psychiatry, disability and tic assessment);
4. CGI – severity by examiner based on video neurological examination;
5. CGI-improvement relative to T = 0 by examiner based on video neurological examination;
6. Counting the amount of jerks during 10 minutes of the psychiatric interview on video;
7. AIMS and PMD scores based on the recorded video during neurological examination;
8. Interview (treatment effective, placebo or BoNT, CGI-S, CGI-I, VAS, adverse events);
9. MRC weakness.

T = 16 months:

1. Neurological examination, recorded on video;
2. Psychiatric interview, recorded on video;
3. Questionnaires (psychiatry, disability and jerk assessment);
4. CGI – severity by examiner based on video neurological examination;
5. CGI-improvement relative to T=0 and T=16 by examiner based on video neurological examination;
6. Counting the amount of jerks during 10 minutes of the psychiatric interview on video;
7. AIMS and PMD scale scores based on the recorded video during neurological examination;
8. Interview (treatment effective, CGI-S, CGI-I, adverse events);
9. MRC weakness.

Intervention

Patients will be randomized to treatment with:

1. 2 times injections with botulinum neurotoxin injections intramuscular with a 3 month interval, or;
2. 2 times injections of placebo intramuscular with a 3 month interval.

Placebo will consist of 0.9% sterile saline. This is a frequently used placebo in studies on botulinum toxin.

After the trial period, all patients will receive 4 injections of BoNT also with 3 month intervals.

For individual patients, the dose of botulinum toxin to be injected will be determined based on the number and type of muscles that are involved in performing the jerk. The muscles that will be injected will be determined by a neurophysiologist experienced in treatment with BoNT according to the pattern of movements visible and according to the involved muscles measured with EMG. The dosage of BoNT depends on the volume of the muscle(s) it is injected in. Guidelines for starting dosages and for adjustments after the first injection for individual muscles exist. These guidelines will be used for the treatment of psychogenic jerks in this study. Because the dosage of BoNT depends on the muscles to be injected, individual patients will receive different dosages. The dilution volume of Dysport will be 500 units in 2.5 cc of 0.9% saline. The maximum dosage per treatment session is 400 Dysport units. BoNT is an accepted and widely-used therapy for dystonia and as such a great deal of experience is available.

Contacts

Public

PO Box 22660
Y.E.M. Dreissen
AMC, D2-136
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5668417

Scientific

PO Box 22660
Y.E.M. Dreissen
AMC, D2-136
Amsterdam 1100 DD
The Netherlands
+31 (0)20 5668417

Eligibility criteria

Inclusion criteria

Patients with at least one invalidating consistent type of psychogenic jerk that is present for 1 year or longer.

Patients with one consistent type of invalidating tremor present for one year or longer .

Exclusion criteria

1. Age < 18 years or > 80 years;
2. Psychogenic jerk of interest present for < 1 year;
3. Treated with BoNT two or more times in the past year;
4. Pregnancy;
5. Coagulation disorders;
6. Insufficient knowledge of Dutch language;
7. Legally incompetent adult;
8. No informed consent.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)
Control:	Placebo

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	07-03-2011

Enrollment: 54
Type: Anticipated

IPD sharing statement

Plan to share IPD: No

Ethics review

Positive opinion
Date: 24-08-2010
Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL2371
NTR-old	NTR2478
Other	MEC Academic Medical Centre : 10/079
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A