

# Gerandomiseerde vergelijkende studie naar verschillen tussen de biologische beschikbaarheid en de effectiviteit van vitamine K drank en vitamine K tabletten.

No registrations found.

<b>Ethical review</b>	Positive opinion
<b>Status</b>	Recruiting
<b>Health condition type</b>	-
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON26078

### Source

Nationaal Trial Register

### Brief title

N/A

### Health condition

Anticoagulants.  
Vitamin K antagonists.  
Vitamin K

Coumarines.  
Vitamine K antagonisten.  
Vitamine K

## Sponsors and support

**Primary sponsor:** Leiden University Medical Center, Department of Thrombosis and Hemostasis

**Source(s) of monetary or material Support:** Dutch Heart Foundation

## Intervention

## Outcome measures

### Primary outcome

1. The difference between vitamin K levels in blood 4 hours after taking vitamin K tablets in comparison with vitamin K solution;
2. The difference in the number of INR-values lower than 2.0 when comparing vitamin K tablets with vitamin K solution in oil after 24 and 48 hours;
3. Difference in the decrease of INRs 24 hours after taking either vitamin K tablets or solution.  
The difference in the number of INRs lower than 2.0 when comparing vitamin K tablets and solution after 24 and 48 hours.

### Secondary outcome

1. The difference between area under the curve measured over 24 hours after taking vitamin K tablets in comparison with vitamin K solution;
2. The difference in increase of the vitamin K levels in the blood 24 hours after taking either vitamin K tablets solution.

## Study description

### Background summary

Vitamin K solution was used for several years to antagonize the treatment with vitamin K antagonists. Vitamin K tablets came into the market as 'vitamin tablets' instead of medicines, but are easier to distribute and do have a longer shelf life. However, they are not shown to be as effective as the oral solution. We perform this study to investigate whether the bioavailability and effectiveness of the tablets and solution are comparable. Patients are therefore recruited from the anticoagulation clinic in Leiden, the Netherlands.

### Study objective

The bioavailability and effectiveness of vitamin K tablets and solution are comparable.

### Study design

Step 1: 0h;2h;4h;5h;6h;8h;10h; 24h

Step 2: 0h; 24h; 48h;

Step 3: 0h; 24h; 48h.

### **Intervention**

Administration of vitamin K tablets or solution.

## **Contacts**

### **Public**

Leiden University Medical Center (LUMC),  
Department of Hematology,  
P.O. Box 9600  
Felix J.M. Meer, van der  
Albinusdreef 2  
Leiden 2300 RC  
The Netherlands  
+31 (0)71 5263901

### **Scientific**

Leiden University Medical Center (LUMC),  
Department of Hematology,  
P.O. Box 9600  
Felix J.M. Meer, van der  
Albinusdreef 2  
Leiden 2300 RC  
The Netherlands  
+31 (0)71 5263901

## **Eligibility criteria**

### **Inclusion criteria**

1. Healthy subjects;
2. Patients being treated with phenprocoumon undergoing an invasive diagnostic or therapeutic procedure for which reducing of the INR is necessary;
3. Patients being treated with phenprocoumon having an INR above 7.0 for which reducing with vitamin K is necessary;

4. Age 18 years or older;
5. Informed consent.

## Exclusion criteria

1. Being treated for liver failure;
2. Pregnancy, pregnancy wish or lactational period;
3. Hemodialysis or peritoneal dialysis;
4. Participation in the self management program;
5. Inability to take care of own medication and/or proven non-compliance with treatment protocols.

## Study design

### Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active

### Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	18-07-2011
Enrollment:	165
Type:	Anticipated

### IPD sharing statement

**Plan to share IPD:** Undecided

## Ethics review

Positive opinion

Date: 18-06-2012

Application type: First submission

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL3353
NTR-old	NTR3485
Other	METC LUMC : P10.177
ISRCTN	ISRCTN wordt niet meer aangevraagd.

## Study results

### Summary results

N/A