Cost-effectiveness of the Demands and Capacities Model based treatment compared to the Lidcombe Programme of early stuttering intervention: Randomised Trial.

No registrations found.

Ethical review	Positive opinion
Status	Recruiting
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON26802

Source Nationaal Trial Register

Health condition

stuttering pre-schoolers

Sponsors and support

Primary sponsor: Dr. Marie-Christine Franken (Erasmus MC, Rotterdam) **Source(s) of monetary or material Support:** ZonMW

Intervention

Outcome measures

Primary outcome

- 1. The percentage of recovered children at 18 months post randomization;
 - 1 Cost-effectiveness of the Demands and Capacities Model based treatment compared ... 23-06-2025

2. The costs for a recovered child.

Secondary outcome

1. The frequency of stuttering outside the clinic at 18 months post randomization. The Health Utility Index, visual analogue scale (Euroqol VAS proxy), Child Health Questionnaire, Children Behaviour Check List.

Study description

Background summary

Stuttering is a frequent problem in three to six year olds: about 5% of children begin to stutter (Bloodstein 1995). The recovery rate four years after onset without intervention is 74% (Yairi & Ambrose 2005). The consensus is that all stuttering children should be treated in the preschool years (Jones et al 2005), because of a higher chance on succesful treatment outcome, presumably so because neural plasticity decreases with age. Also, it is nog yet possible to predict for an individual case who will or will not recover from stuttering. In the Netherlands, the Demands and Capacities Model based treatment is the standard (DCM, Starkweather et al 1990). This treatment uses an indirect approach. The Lidcombe Programme (LP) is the Australian standard for treating pre-school children who stutter. This is a direct treatment, because it aims to increase fluent and decrease stuttered speech. Evidence suggests that both treatments are effective. A RCT with the LP showed that at 9 months after randomization, children who were treated with the LP were stuttering less than children in the no treatment arm (Jones et al 2005). Efficacy of the DCM has been shown in a randomized pilot trial in which the effects of DCM and LP treatment after 12 weeks were compared to each other (Franken et al 2005). To improve the evidence basis underlying stuttertherapies, a cost-effective evaluation of stuttering therapy is relevant. The objective of this study is to determine the relative effectiveness, cost-effectiveness and cost-utility of the Dutch standard for treating stuttering in pre-school children (DCM) compared to the Australian standard (LP).

Outcomes will be analyzed at baseline and at 3, 6, 12 and 18 months post randomization.

Study objective

The Lidcombe Programme for early stuttering intervention is more cost-effective than the Demands and Capacities Model based treatment.

Intervention

Demands and Capacities Model based treatment versus Lidcombe Programme

Contacts

Public

Erasmus MC Rotterdam, afdeling iBMG, Postbus 1738 C. Koedoot Rotterdam 3000 DR The Netherlands +31 10-4088617 **Scientific** Erasmus MC Rotterdam, afdeling iBMG, Postbus 1738 C. Koedoot Rotterdam 3000 DR The Netherlands +31 10-4088617

Eligibility criteria

Inclusion criteria

- 1. Age 3.0-6.3;
- 2. Frequency of stuttering at least 3%;
- 3. Parent and one therapist agree the child stutters;
- 4. Parent rating of stuttering severity on an 8-point scale of at least 2;
- 5. Proficiency in Dutch for children and parents.

Exclusion criteria

- 1. Onset of stuttering within 6 months before recruitment;
- 2. Treatment for stuttering during the previous 12 months;
- 3. Diagnosed language disorder;
- 4. Neurologic, emotional, cognitive, behavioral or autism spectrum disorder.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Masking:	Single blinded (masking used)
Control:	Active

Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	01-08-2007
Enrollment:	196
Туре:	Anticipated

Ethics review

Positive opinion		
Date:	23-07-2007	
Application type:	First submission	

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL995
NTR-old	NTR1024

4 - Cost-effectiveness of the Demands and Capacities Model based treatment compared ... 23-06-2025

Register		
Other		
ISRCTN		

Study results

Summary results

Franken, M.C.J., Kielstra-van der Schalk, C.J. & Boelens, H. (2005). Experimental treatment of early stuttering: a preliminary study. Journal of Fluency Disorders, 30 (3), 189-199.

ID :

ISRCTN24362190

Jones, M., Onslow, M. Packman, A., Williams, S., Ormond, T., Schwarz, I., Gebski, V. (2005). Randomised controlled trial of the Lidcombe programme of early stuttering intervention. British Medical Journal, 331, 659-661.

Gottwald, S.R. & Starkweather, C,W. (1999). Stuttering prevention and early intervention: a multiprocess approach. in; M. Onslow & A. Packman (eds.) The handbook of early stuttering intervention, 53-82. San Diego/London: Singular Publishing Group.