

Neural plasticity, hearing loss and tinnitus: changes in tonotopic maps of the auditory cortex

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Observational non invasive

Summary

ID

NL-OMON27684

Source

Nationaal Trial Register

Health condition

hearing loss
tinnitus
tonotopy
auditory cortex

Sponsors and support

Primary sponsor: University Medical Center Groningen

Source(s) of monetary or material Support: University Medical Center Groningen.
American Tinnitus Association

Intervention

Outcome measures

Primary outcome

Measures of tonotopic map reorganisation.

Secondary outcome

Relation of hearing loss and degree of reorganisation. Two types of hearing loss are going to be studied: steep-sloping hearing loss and gradual sloping hearing loss.

Study description

Background summary

With an increased life expectancy and an ageing population, age-related dysfunctions are becoming more prevalent. The most widespread of all sensory impairments in an ageing population is hearing loss, which is characterized by a loss of sensitivity of the peripheral hearing organ, the inner ear. This peripheral hearing loss is associated with less sensory input available to the brain. Animal research has shown that peripheral hearing loss may cause widespread plastic changes in the brain. Peripheral hearing loss is often associated with tinnitus: about 30% of the people with hearing loss also develop tinnitus. It has been suggested previously that tinnitus and tonotopic reorganisation are causally related. Such cortical reorganisation presumably contributes to the impaired communication skills experienced by humans with hearing loss. With an increasing demand for active participation of the elderly in society, it is crucially important to understand the neurobiological consequences of hearing loss and tinnitus.

The current study employs functional magnetic resonance imaging (fMRI) to investigate the relation between peripheral hearing loss, tinnitus and cortical reorganization. In particular, it aims to map the topographic representation of sound frequency, referred to as tonotopic maps, and how these change as a consequence of tinnitus and hearing loss. It also aims to investigate whether the type of hearing loss is related to the degree of reorganisation.

Study design

No longitudinal design; just one time-point

Intervention

fMRI measurements while listening to sound fragments.

Contacts

Public

C.P. Lanting
Faculteit Medische Wetenschappen/UMCG, Keel- Neus- en Oorheelkunde
Hanzeplein 1
Groningen 9713 GZ
The Netherlands
050 363 5111

Scientific

C.P. Lanting
Faculteit Medische Wetenschappen/UMCG, Keel- Neus- en Oorheelkunde
Hanzeplein 1
Groningen 9713 GZ
The Netherlands
050 363 5111

Eligibility criteria

Inclusion criteria

- No hearing loss or reported tinnitus (controls, n=40)
- Hearing loss (Hearing loss group, n=40)
- Tinnitus (Hearing loss and tinnitus group, n=40)
- < 30 dB difference between both ears for all the standard audiometric frequencies
- Adults (18 - 75 yrs.)
- No contraindications for MRI

Exclusion criteria

Non-compliance with inclusion criteria

Study design

Design

Study type: Observational non invasive

Intervention model:	Other
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-01-2015
Enrollment:	120
Type:	Anticipated

Ethics review

Positive opinion	
Date:	24-09-2014
Application type:	First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 40455
Bron: ToetsingOnline
Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL4642
NTR-old	NTR4811
CCMO	NL44470.042.13
OMON	NL-OMON40455

Study results