

# The association between a drop in systolic blood pressure from the individual reference value and in-hospital mortality in ED patients.

No registrations found.

<b>Ethical review</b>	Positive opinion
<b>Status</b>	Recruiting
<b>Health condition type</b>	-
<b>Study type</b>	Observational non invasive

## Summary

### ID

NL-OMON27709

### Source

Nationaal Trial Register

### Brief title

DeltaSBP

### Health condition

Presentation in the ED

## Sponsors and support

**Primary sponsor:** none

**Source(s) of monetary or material Support:** Maxima Medical Centre

## Intervention

## Outcome measures

### Primary outcome

in-hospital mortality

## Secondary outcome

ICU admission

## Study description

### Background summary

Risk stratification of ED patients is important for appropriate initial treatment and disposition to a ward or intensive care unit (ICU). Clinical deterioration of patients often starts in the prehospital setting and only when patients start to feel too sick they will visit the GP or Emergency Department (ED). In a previous study we showed that the odds for in-hospital mortality increased approximately linearly with decreasing systolic blood pressure (SBP) starting at 140mmHg for unselected ED patients. Thus, on an average cohort level, prognosis of ED patients deteriorates already with SBPs below 140 mmHg. This may be explained by a relatively small number of ED patients who had a much higher SBP than 140 mmHg before they presented to the ED, as hypertension is a very common chronic disease in Western countries. If so, not only the initial SBP in the ED is relevant for prognosis but the absolute or relative reduction of the SBP which was normal for the individual patient before the ED presentation, i.e. it is important to know the baseline SBP of the individual patient. However, this information may often not be available in the ED as electronic patient records (EPR) are not routinely shared between general practitioners and hospitals in the Netherlands, due to privacy laws. In addition, in the outpatient clinics, blood pressure is not routinely measured. Before we can investigate the association between the absolute or relative reduction of SBP from the individual baseline SBP, we need to know in how many patients baseline SBPs are available. If indeed many baseline values are unavailable while the reduction of SBP is associated for the prognosis of the individual ED patient, this may plead for the need of a shared Electronic Patient Record (EPR) with the necessary clinical information, such as baseline vital signs. More importantly, if we find an association between a reduction of SBP from the individual baseline SBP and in-hospital mortality, a larger prospective cohort study would be indicated, which would then help to improve risk stratification of individual ED patients.

The aim of the present pilot study is therefore two-fold:

#### Objectives

1. To assess how often individual baseline values for SBP are known in ED patients.
2. To assess whether a change in SBP from the individual baseline values for SBP are associated with in-hospital mortality.

### Study objective

A low delta SBP is associated with an increased risk of mortality

### Study design

We study overall in-hospital mortality or ICU admission from the ED. After discharge we don't follow-up on patients.

## **Intervention**

no

## **Contacts**

### **Public**

Leiden Universitair Medisch Centrum  
Bart Candel

0644665616

### **Scientific**

Leiden Universitair Medisch Centrum  
Bart Candel

0644665616

## **Eligibility criteria**

### **Inclusion criteria**

Patients >70 years with ED visit and admission

### **Exclusion criteria**

triage category blue/green

## **Study design**

### **Design**

Study type: Observational non invasive

Intervention model: Factorial

Allocation:	Non controlled trial
Masking:	Open (masking not used)
Control:	Active

## Recruitment

NL	
Recruitment status:	Recruiting
Start date (anticipated):	03-11-2020
Enrollment:	220
Type:	Anticipated

## IPD sharing statement

**Plan to share IPD:** No

## Ethics review

Positive opinion	
Date:	03-11-2020
Application type:	First submission

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
NTR-new	NL9029
Other	METC Máxima MC : N20.052

## Study results