

Prospectief gerandomiseerd multicenter vergelijkend onderzoek tussen het gebruik van rubberband ligatie en arterie ligatie van de hemorroïdaal arteriën voor de behandeling van graad II en III hemorroïden.

No registrations found.

Ethical review	Positive opinion
Status	Pending
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON27930

Source

Nationaal Trial Register

Brief title

HEMO

Health condition

Hemorrhoids, piles, THD, HAL, rubberband ligation.

Hemorroïden, aambeien, THD, HAL, Rubberband ligatie

Sponsors and support

Primary sponsor: St. Antonius Ziekenhuis, Nieuwegein, Netherlands

Source(s) of monetary or material Support: fund = initiator = sponsor

Intervention

Outcome measures

Primary outcome

Primary endpoint is defined as a successful treatment out of patient perspective. A treatment is considered successful if the number one complaints has subsided.

Secondary outcome

Secondary endpoints are:

1. Cost effectivity;
2. Number of treatments necessary;
3. Number of clinical visits;
4. Number of days not able to work;
5. Number of days of hospitalisation;
6. Postoperative pain;
7. Amount of painmedication used post operative;
8. Post operative complication in the first week after the procedure and during 6 months of follow-up;
9. Number of re-interventions.

Study description

Background summary

In daily practice there are many treatment modalities for hemorrhoids available. Since many years the rubberband ligation is widely used. Recently there are also positive results reported of the artery ligation procedure. This research aims to determine best practice in the treatment of grade II and III hemorrhoids.

The aim of this study is to determine which of the two common treatment modalities of hemorrhoids known as rubberband ligation and artery ligation procedure is best for the

treatment of grade II or III hemorrhoids.

Study objective

In daily practice there are many treatment modalities for hemorrhoids available. Since many years the rubberband ligation is widely used. Recently there are also positive results reported of the artery ligation procedure. This research aims to determine best practice in the treatment of grade II and III hemorrhoids.

Study design

Follow-up at 6 weeks, and 6 months.

Intervention

Patients are either treated with the rubberband ligation procedure or with the artery ligation procedure.

Contacts

Public

Postbus 2500
Jaap-Peter Schuurman
Nieuwegein 3430 EM
The Netherlands
+31 (0)88 320 30 00

Scientific

Postbus 2500
Jaap-Peter Schuurman
Nieuwegein 3430 EM
The Netherlands
+31 (0)88 320 30 00

Eligibility criteria

Inclusion criteria

1. By means of proctoscopy, physical examination or sigmoidoscopy proved grade II or III hemorrhoids;
2. No other explanation for rectal bloodloss;

3. Age between 18 and 70 years;
4. Patient understands the Dutch language;
5. Patient is able to decide;
6. Informed consent.

Exclusion criteria

1. Grade I or IV hemorrhoids;
2. Asymptomatic hemorrhoids;
3. Other proctologic disorders;
4. Coagulopathy;
5. Use of coumarine derivatives;
6. Anal surgery in history.

Study design

Design

Study type:	Interventional
Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	N/A , unknown

Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	01-01-2010
Enrollment:	200
Type:	Anticipated

Ethics review

Positive opinion

Date: 18-11-2009

Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL1999
NTR-old	NTR2116
Other	CCMO : ABR26912 / NL26912.100.09
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Study results

Summary results

N/A