

CasWhey study: Casein protein versus Whey protein

No registrations found.

Ethical review	Positive opinion
Status	Recruitment stopped
Health condition type	-
Study type	Interventional

Summary

ID

NL-OMON28888

Source

Nationaal Trial Register

Brief title

CasWhey study

Health condition

Amino acid absorption after gastric bypass surgery

Satiety

Secretion of anorexigenic hormones

Sponsors and support

Primary sponsor: Medical Center Leeuwarden

Source(s) of monetary or material Support: 1. University of Groningen/Campus Fryslân
2. Medical Centre Leeuwarden
3. University Medical Centre Groningen / Department of Endocrinology

Intervention

Outcome measures

Primary outcome

- To assess and compare the net incremental area under the curve of satiety-related plasma

amino acids and derivatives in response to lactase-containing milk supplemented with either casein or whey protein in a randomized controlled cross over study in subjects 12-18 months after primary gastric bypass surgery.

Secondary outcome

- To assess and compare the effect of the ingestion of both protein supplements in lactase-containing milk on plasma concentrations of all amino acids and derivatives, plasma concentrations of anorexigenic gastrointestinal hormones PYY and GLP-1, satiety, hunger, dumping symptoms, heart rate, blood pressure, glucose metabolism and ad libitum food intake after completion of the test meal in a randomized controlled cross over study in subjects 12-18 months after primary gastric bypass surgery.
- To assess the self-reported dietary patterns, macronutrient consumption, energy intake, eating behaviour and physical activity measured by questionnaires in subjects 12-18 months after primary gastric bypass surgery.

Study description

Background summary

Rationale: Dietary protein could be of potential great importance to improve postoperative outcomes after bariatric surgery given its effects in the non-surgical population on preservation of fat free mass, induction of satiety and prevention of protein malnutrition. And while the quantity of dietary protein is the main focus of interest in many studies, the amino acid composition of the ingested proteins may also be of importance. Our systematic review on the effect of dietary protein after gastric bypass surgery concluded there might be an essential beneficial role, but due to limited amount of literature and the diversity in main outcomes solid evidence could not be provided and more high quality research is needed. Results from our recent study showed that several plasma concentrations of amino acids were elevated in gastric bypass patients after a mixed meal tolerance test in those who experience more satiety and in those with higher concentrations of satiety-related (anorexigenic) hormones in 3.5 hours after ingestion of the test meal. This study points in the direction that a postprandial plasma amino acid profile comparable to casein protein is associated with both increased satiety and increased concentrations of satiety-related gastrointestinal hormones. However, in healthy and obese subjects whey protein induced more effect on satiety, active GLP-1 and insulin compared to casein protein. No studies with different types of proteins have been performed in gastric bypass patients.

Objective: To investigate the effect of a test meal supplemented with casein protein in gastric bypass patients on concentrations of satiety-related plasma amino acids and satiety-related hormones and feelings of postprandial satiety and hunger compared to a test meal supplemented with whey protein.

Study design: This is a pilot study with a blinded randomized cross-over design. The recruitment will start in November 2017 and continue by approaching potential candidates consecutively.

Study population: In order to be eligible for participation in this study, subjects are women aged 18 to 65 years during their primary gastric bypass surgery 12-18 months ago at the Centre of Obesity Netherlands at the Medical Centre Leeuwarden. In total 18 subjects will be evaluated.

Intervention: During two study visits the subjects will receive 200mL lactase-containing milk with either 15 g casein protein or 15 g whey protein as supplement in a blinded randomized order.

Main study endpoint: The difference in net incremental area under the curves of the plasma concentrations of satiety-related amino acids and derivatives after ingestion of the two test meals.

Study objective

-

Study design

18 subjects will receive either casein protein or whey protein in a randomized order during two visits with a period of minimally 1 week and maximally 3 weeks between them on the same day of the week.

Each visit blood will be withdrawn through an intravenous catheter on 9 time points. An overview of the tubes is presented in Table 2. In total, 193.5 (21.5mL x 9 time points) mL blood will be withdrawn for the analyses per visit.

Intervention

During two study visits the subjects will receive 200mL lactase-containing milk with either 15 g casein protein or 15 g whey protein as supplement in a blinded randomized order.

Contacts

Public

Medical Centre Leeuwarden

Merel van den Broek
Henri Dunantweg 2

Leeuwarden 8934 AD
The Netherlands
+3158-2861940

Scientific

Medical Centre Leeuwarden

Merel van den Broek
Henri Dunantweg 2

Leeuwarden 8934 AD
The Netherlands
+3158-2861940

Eligibility criteria

Inclusion criteria

Female

Gastric bypass surgery at age 18-65 years

Primary gastric bypass between 12 and 18 months ago

Exclusion criteria

Follow-up is not feasible

Known allergies or intolerance to ingredients used in the intervention product

- o Milk

- o Soy

Medicine usage known to influence dietary uptake and/or appetite

- o Antidepressants

- o Antiemetics
- o Laxatives
- o Levodopa
- o Lipid-lowering medication
- o Opiates
- o Prokinetics
- o Systemic corticosteroids

Pregnancy

Hypo- or hyperthyroidism

Known malabsorption syndrome

Known heart failure

Known renal insufficiency or failure

Known severe hepatic disease

Known severe symptomatic post gastric bypass hypoglycaemia

Active Diabetes Mellitus or remission

Former gastric operations

Study design

Design

Study type:	Interventional
Intervention model:	Crossover
Allocation:	Randomized controlled trial
Masking:	Single blinded (masking used)
Control:	N/A , unknown

Recruitment

NL
Recruitment status: Recruitment stopped
Start date (anticipated): 03-01-2018
Enrollment: 18
Type: Actual

IPD sharing statement

Plan to share IPD: Undecided

Ethics review

Positive opinion
Date: 31-07-2017
Application type: First submission

Study registrations

Followed up by the following (possibly more current) registration

ID: 44185
Bron: ToetsingOnline
Titel:

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
NTR-new	NL6459
NTR-old	NTR6637
CCMO	NL61997.099.17
OMON	NL-OMON44185

Study results