

# The emotional response in Parkinson`s disease before and after bilateral STN stimulation

Published: 04-12-2006

Last updated: 14-05-2024

To gain knowledge about the etiology of emotional changes after bilateral STN stimulation. And also to find out that the research design, where neuropsychological and psysiological aspects are combined, is useful in studying outcomes of STN...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Pending
<b>Health condition type</b>	Movement disorders (incl parkinsonism)
<b>Study type</b>	Observational non invasive

## Summary

### ID

NL-OMON30351

### Source

ToetsingOnline

### Brief title

Emotional respons and STN

### Condition

- Movement disorders (incl parkinsonism)

### Synonym

Parkinson movementdisorder

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Academisch Medisch Centrum

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** Emotional response Parkinson STN

## Outcome measures

### Primary outcome

auditive startle reflex (latency, duration and peak of the amplitude EMG activity per muscle)

tactile startle reflex (latency, duration and peak of the amplitude EMG activity per muscle)

Psychogalvanic respons (difference between the minimum and maximum activity in a period of four seconds after stimulus onset)

Affective reports on arousal and valence (Self Assessment Manikin)

### Secondary outcome

General cognitive functioning (Mattis DRS)

Executive functions (letterfluency, Trail Making Test en Stroop)

Depression and/or anxiety (HADS)

Mood states (PANAS/POMS)

Motor symptoms (UPDRS part 3)

## Study description

### Background summary

Patients with advanced Parkinson`s disease often have response fluctuations and dyskinesias. Bilateral subthalamic nucleus (STN) stimulation is a possibility for reducing motor symptoms. However, negative outcomes have been reported, like the presence of emotional changes.

A possible explanation is alternation of the emotional response after STN stimulation. Lang, Bradley and Cuthbert (1990) define emotions as action dispositions, which are organized around two dimensions: valence and arousal. Emotional response can be measured by affective reports and physiological activity, in particular the startle reflex and psychogalvanic response (PGR). Patients with Parkinson`s disease have a reduced physiological response to negative stimuli. Affective reports were similar to reports of controls (Miller, 2004). Research of the emotional response after bilateral STN stimulation has not been reported yet. Therefore the research question of this study is: Is there a change of emotional response in Parkinson`s disease after bilateral STN stimulation?

### **Study objective**

To gain knowledge about the etiology of emotional changes after bilateral STN stimulation. And also to find out that the research design, where neuropsychological and psysiological aspects are combined, is useful in studying outcomes of STN stimulation.

### **Study design**

The study consists of a controlled, repeated measures design. Patients with Parkinson`s disease who will undergo an STN stimulation will be examined shortly before and two months after the surgery. Patients with Parkinson`s disease who do not undergo surgery will be asked as controls. The auditory and tactile startle reflex will be studied before and after surgery. Moreover pleasant, unpleasant and neutral pictures will be showed. At the same time the PGR is measured. Also, at each picture affective reports on valence and arousal will be determined. Finally, tests concerning general cognitive functioning, executive function, anxiety, depression, mood states and motor symptoms are administered. Each evaluation will take at most two hours.

### **Study burden and risks**

Each evaluation will take at most two hours. First light stress is caused by auditory and tactile stimuli. This procedure contains no risks for the patient. Moreover tests concerning general cognitive functioning, executive function, anxiety, depression, mood states and motor symptoms are also administered.

## Contacts

### Public

Academisch Medisch Centrum

postbus 22660  
1100 DD Amsterdam  
Nederland

### Scientific

Academisch Medisch Centrum

postbus 22660  
1100 DD Amsterdam  
Nederland

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

- disease duration et least 5 years or more
- experimental condition: patiënts who undergo an STN stimulation

### Exclusion criteria

- dementia (Mattis DRS < 120)
- other serious central nervous system diseases or other diseases.

## Study design

### Design

Study type:	Observational non invasive
Intervention model:	Other
Allocation:	Non-randomized controlled trial
Masking:	Open (masking not used)

**Primary purpose:** Basic science

### Recruitment

NL	
Recruitment status:	Pending
Start date (anticipated):	20-10-2006
Enrollment:	10
Type:	Anticipated

## Ethics review

Approved WMO	
Application type:	First submission
Review commission:	METC Amsterdam UMC

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

### Register

CCMO

### ID

NL14797.018.06