

Effect of preoperative pelvic floor physiotherapy versus standard care on incontinence in men undergoing radical laparoscopic prostatectomy

Published: 21-09-2007

Last updated: 08-05-2024

NA

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Urolithiases
Study type	Observational invasive

Summary

ID

NL-OMON30981

Source

ToetsingOnline

Brief title

Effect of preoperative pelvic floor physiotherapy

Condition

- Urolithiases

Synonym

urinary incontinence

Research involving

Human

Sponsors and support

Primary sponsor: Leids Universitair Medisch Centrum

Source(s) of monetary or material Support: Ministerie van OC&W

Intervention

Keyword: laparoscopic prostatectomy, Pelvic floor

Outcome measures

Primary outcome

NA

Secondary outcome

NA

Study description

Background summary

NA

Study objective

NA

Study design

NA

Study burden and risks

NA

Contacts

Public

Leids Universitair Medisch Centrum

Albinusdreef 2
2333 ZA Leiden
NL
Scientific
Leids Universitair Medisch Centrum

Albinusdreef 2
2333 ZA Leiden
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

prostate cancer

Exclusion criteria

neurological diseases

Study design

Design

Study type: Observational invasive

Intervention model: Parallel

Allocation:	Randomized controlled trial
Masking:	Double blinded (masking used)

Primary purpose: Diagnostic

Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	15-11-2007
Enrollment:	250
Type:	Actual

Ethics review

Approved WMO	
Application type:	First submission
Review commission:	METC Leids Universitair Medisch Centrum (Leiden)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL17911.058.07