

# European multi centre trial to compare the effectiveness of two different phototherapy narrow band UVB protocols in vitiligo

Published: 27-05-2008

Last updated: 07-05-2024

The primary objective of this study is to compare the effectiveness of two different phototherapy protocols (continuous treatment for 6 months, or 2 continuous treatments for 2 months separated by one month interruption (2-1-2-1), with a total...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Pending
<b>Health condition type</b>	Pigmentation disorders
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON32305

### Source

ToetsingOnline

### Brief title

MCT nb UVB

### Condition

- Pigmentation disorders

### Synonym

vitiligo, white spot disease

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Academisch Medisch Centrum

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** MCT, nb UVB, phototherapy, vitiligo

## Outcome measures

### Primary outcome

Colour measurement by visual assessment and reflectance spectroscopy, patient\*s satisfaction and visual assessment of side effects.

### Secondary outcome

not applicable

## Study description

### Background summary

Narrow band ultraviolet B (nb UVB) phototherapy is the first choice for the treatment of active vitiligo (newly developed depigmented macules in the last 3 months). Phototherapy requires quite long periods (6 months \* 1 year) before repigmentation of vitiligo takes place. Side effect of phototherapeutic treatment is represented by the cumulative long term effects of ultraviolet rays: premature ageing of the skin (wrinkles, spots etc.) and the appearance of skin cancer/neoplasia. Our aim is to get the best results from phototherapy whilst limiting its length.

### Study objective

The primary objective of this study is to compare the effectiveness of two different phototherapy protocols (continuous treatment for 6 months, or 2 continuous treatments for 2 months separated by one month interruption (2-1-2-1), with a total length of 6 months as well). Our aim is to verify whether interrupting a cycle of phototherapy is useful or not. This interruption might make the ultraviolet rays\* stimulus on cells producing melanin more effective, and may also reduce the long term damage caused by ultraviolet rays.

### Study design

Prospective single blinded randomised controlled multi centre study.

## **Intervention**

nb UVB phototherapy

## **Study burden and risks**

Subjects participating in the study will be asked to visit the SNIP 5 times. The time investment will be 15 minutes per visit. No invasive procedures will be performed. The objective colour measurement involves a handheld device producing a harmless flash of light. Known side effects of the nb UVB phototherapy are redness, pruritus, xerosis cutis, burning sensation and conjunctivitis. All together the burden due to the study is moderately low and the risk for local side effects is low. Systemic side effects are not associated with the involved treatment.

## **Contacts**

### **Public**

Academisch Medisch Centrum

Meibergdreef 35  
1105AZ, Amsterdam  
Nederland

### **Scientific**

Academisch Medisch Centrum

Meibergdreef 35  
1105AZ, Amsterdam  
Nederland

## **Trial sites**

### **Listed location countries**

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

Active vitiligo (newly developed depigmented macules in the last 3 months)

Subjects attending the outpatient department of the SNIP;

Aged between 18-60 years;

Subject is willing and able to give written informed consent.

### Exclusion criteria

Subjects not competent to understand what is involved.

Skin photo type I

Previous treatment with any kind of phototherapy in the last 6 months

Acral vitiligo (only hands and feet are affected)

Lesion(s) suspicious for malignancy

Pregnancy

High exposure to sunlight (vacation in southern countries) or UV light (UVA or UVB)

## Study design

### Design

**Study type:** Interventional

Masking: Single blinded (masking used)

Control: Uncontrolled

Primary purpose: Treatment

### Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-03-2008

Enrollment: 20

Type:

Anticipated

## Ethics review

Approved WMO

Application type:

First submission

Review commission:

METC Amsterdam UMC

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

**Register**

CCMO

**ID**

NL21475.018.08