

# An open phase 1, dose-escalating, clinical trial on the safety of a new liposomal adjuvant system, CAF01, when given with the tuberculosis subunit vaccine Ag85B-ESAT-6 as two injections with two months interval to healthy adult volunteers.

Published: 08-06-2009

Last updated: 06-05-2024

**Primary:** The primary objective is to evaluate the safety profile of an adjuvated Tb subunit vaccine (CAF01) administered in 50 µg Ag85B-Esat-6 alone, 50 µg Ag85B-Esat-6 alone with three escalating CAF01 dose levels, to four groups of healthy...

<b>Ethical review</b>	-
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Mycobacterial infectious disorders
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON33959

### Source

ToetsingOnline

### Brief title

ACAF01-01

### Condition

- Mycobacterial infectious disorders

### Synonym

Prevention, Tuberculosis

## **Research involving**

Human

## **Sponsors and support**

**Primary sponsor:** Statens Serum Institut

**Source(s) of monetary or material Support:** SSI (Statens Serum Institut)

## **Intervention**

**Keyword:** ADJUVATED, HEALTHY VOLUNTEERS, TUBERCULOSIS, VACCINE

## **Outcome measures**

### **Primary outcome**

Safety variables:

- Adverse reactions with onset between the first vaccination and 24 weeks after the second vaccination either identified during medical examinations, as diary records, or during safety interviews

- Changes from baseline in safety laboratory test values 1 day, 7 days and 6 weeks after the first and second vaccinations and 24 weeks after the second vaccination.

### **Secondary outcome**

Immunogenicity variables

- Changes from baseline in Ag85B-Esat-6 induced IFN-gamma release from PBMC's measured by ELISA, 1 week and 6 weeks after the first and second vaccinations and 24 and 44 weeks after the second vaccination.

- Changes from baseline in Ag85B-ESAT6 IFN-gamma spot forming cells measured by ELISPOT, 1 week and 6 weeks after the first and second vaccinations and 24 and 44 weeks after the second vaccination.

- Changes from baseline in serum Ig-G antibodies against Ag85B-ESAT-6, 6 weeks after the first and second vaccinations and 24 and 44 weeks after the second vaccination.

## Study description

### Background summary

Title:

A safety and immunogenicity phase 1 trial with an adjuvated TB subunit vaccine (Ag85B - Esat-6 + CAF01) administered in PPD positive volunteers at 0 and 2 months.

Background:

Tuberculosis (TB) is caused by *Mycobacterium tuberculosis* (MT), an intracellular pathogen. One third of the world's population is infected with TB, 8-10 million suffer from TB-disease and 2-3 million die annually. Currently the only available vaccine against TB is BCG. BCG protects against severe childhood forms of TB. However, the protective efficacy in adult pulmonary tuberculosis varies considerably, from 85% to 0%. A new improved second generation TB vaccine is therefore urgently needed.

### Study objective

Primary:

The primary objective is to evaluate the safety profile of an adjuvated Tb subunit vaccine (CAF01) administered in 50 µg Ag85B-Esat-6 alone, 50 µg Ag85B-Esat-6 alone with three escalating CAF01 dose levels, to four groups of healthy volunteers at 0 and 2 months, injecting two doses.

Secondary:

The secondary objective is to determine the immunogenicity profile of an

adjuvant TB subunit vaccine (CAF01) administered in 50 µg Ag85B-Esat-6 alone, 50 µg Ag85B-Esat-6 alone with three escalating CAF01 dose levels, to four groups of healthy volunteers at 0 and 2 months, injecting two doses.

## Study design

A single-centre, open, phase 1 trial with a fixed antigen but a variable doses of adjuvant, including four groups of volunteers. In total 37 volunteers, vaccinated twice, 0 and two months. A data Safety Monitoring Board (DSMB) decided if it is safe to continue with the second vaccination.

## Intervention

Two vaccinations over a two month period, intramuscularly into deltoid muscle.

## Study burden and risks

Twice a vaccination. Ten times blood withdrawal, Mantoux-skin-test.

Risks: anafylactic shock caused by the vaccine.

## Contacts

### Public

Statens Serum Institut

Artillerivej 5  
DK-2300 Copenhagen S  
DK

### Scientific

Statens Serum Institut

Artillerivej 5  
DK-2300 Copenhagen S  
DK

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

Female and male (18-55 years)

Healthy

### Exclusion criteria

History of tuberculosis or known exposure

Positive Tuberculin Skin Test and/or positive Quantiferon

BCG vaccination

Thyroid dysfunction

Disease affecting the lymphoid organs

ANA-titer, HBV, HCV, HIV

C-reactive protein level >50 mg/l

Live vaccine vaccination within 3 months before the first vaccination

Intake of another clinical trial product/vaccine within 3 months from the first vaccination or participation in previous clinical trials with the Ag85B-Esat-6 antigen.

## Study design

### Design

**Study type:** Interventional

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Prevention

### Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 16-11-2009

Enrollment: 37  
Type: Actual

## Ethics review

Approved WMO  
Date: 06-11-2012  
Application type: Amendment  
Review commission: CCMO: Centrale Commissie Mensgebonden Onderzoek (Den Haag)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
EudraCT	EUCTR2008-006003-23-NL
CCMO	NL26270.000.09