

# PTH after total thyroidectomy, a predictive factor for long-term hypoparathyroidism?

Published: 11-02-2014

Last updated: 24-04-2024

The aim of our study is to investigate whether PTH levels measured between 4-24 hours after surgery can accurately identify patients with permanent hypoparathyroidism after thyroid surgery who need long-term calcium supplementation.

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Pending
<b>Health condition type</b>	Thyroid gland disorders
<b>Study type</b>	Observational non invasive

## Summary

### ID

NL-OMON38463

### Source

ToetsingOnline

### Brief title

PTH after total thyroidectomy

### Condition

- Thyroid gland disorders

### Synonym

hypoparathyroidism

### Research involving

Human

### Sponsors and support

**Primary sponsor:** Erasmus MC, Universitair Medisch Centrum Rotterdam

**Source(s) of monetary or material Support:** Ministerie van OC&W

## Intervention

**Keyword:** calcium supplementation, hypoparathyroidism, parathyroidhormone, Thyroidectomy

## Outcome measures

### Primary outcome

Can we use PTH levels obtained directly after surgery to predict which patient will remain hypocalcaemic?

### Secondary outcome

Not applicable

## Study description

### Background summary

Total thyroidectomy is the operation of choice for many patients with thyroid carcinoma, and sometimes with Graves\* disease or multinodular goitre. Postoperative hypoparathyroidism, following total thyroidectomy is a common iatrogenic complication leading, among other factors, to hypocalcaemia. Up to 55-60% of the patients develop serum calcium levels below 2.00 mmol/L on the first day after surgery. Therefore it is routine practice to monitor patients clinically for signs and symptoms of hypocalcaemia as well as perform regular measurements of serum calcium levels. Recent studies (2,3) have shown that a serum parathyroid hormone (PTH) concentration obtained 4-24 hours postoperatively is the most significant factor to predict postoperative hypocalcaemia. In most patients with post-thyroidectomy hypocalcaemia, parathyroid function (if any glands are left) restores within several weeks. However, up to 5-10% of the patients may develop persistent hypoparathyroidism and require lifelong calcium supplementation. In a study by Sitges-Serra et al.<sup>1</sup> evaluating 442 patients after total thyroidectomy, 222 patients had postoperative hypocalcaemia. Of them, 40% (80/222 patients) still had hypocalcaemia one month after surgery. PTH levels were also measured one month after surgery: 90% (40/44 patients) of the patients with low PTH levels recovered to normal parathyroid function within one year compared with 62% of the patients with undetectable parathyroid levels (21/34 patients). Logistic regression analysis identified undetectable PTH levels one month after surgery to predict permanent failure of parathyroid function. However, PTH levels directly after surgery were not measured. We aim to identify a predictive

factor for persistent hypoparathyroidism after a total (or completion) thyroidectomy that can be obtained directly after surgery. This would not only be valuable for planning outpatient appointments, but would also provide more certainty for the patient. In addition calcium supplementation can become more patient tailored at the outpatient department, aiming to reduce the negative consequences of hypocalcaemia, unnecessary visits and laboratory investigations. Therefore the aim of our study is to investigate whether PTH levels measured between 4-24,3 hours after surgery will identify patients who need long-term calcium supplementation.

## **Study objective**

The aim of our study is to investigate whether PTH levels measured between 4-24 hours after surgery can accurately identify patients with permanent hypoparathyroidism after thyroid surgery who need long-term calcium supplementation.

## **Study design**

This extra protocol applies to all patients who have given informed consent.

1. Prior to surgery (outpatient department), (Day -1)
  - a. Check informed consent and contact with study coordinator
  - b. Lab (t=0): Routine laboratory screening, including: calcium/ phosphate/albumin/vitamin D including: PTH for the study
2. Day of surgery (Day = 0)
  - a. Lab (t=1 / 18:00): Routine laboratory screening (calcium, phosphate, albumin, vitamin D), including: PTH for the study
3. Day after surgery (Day = +1)
  - a. Lab (t=2 / 08:00): Routine laboratory screening (calcium, phosphate, albumin), including: PTH for the study
4. 1-2 weeks after surgery
  - a. Lab (t=3): Routine laboratory screening (calcium, phosphate, albumin), including: PTH for the study
5. 1 year after surgery
  - a. Lab (t=4): Routine laboratory screening (calcium, phosphate, albumin), including: PTH for the study

Interpretation of laboratory findings and calcium supplementation are based on the existing protocol (Attachment 1) which is not changed for the purpose of the study.

## **Study burden and risks**

not applicable.

## Contacts

### Public

Erasmus MC, Universitair Medisch Centrum Rotterdam

s'-gravendijkwal 213  
Rotterdam 3015 CE  
NL

### Scientific

Erasmus MC, Universitair Medisch Centrum Rotterdam

s'-gravendijkwal 213  
Rotterdam 3015 CE  
NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

### Inclusion criteria

All patients scheduled for a total thyroidectomy (or subtotal if the isthmus is to remain in situ) or completion thyroidectomy are to be included in this study.

### Exclusion criteria

Patients undergoing concomitant lymph node dissection may also be included in the study.

- o Age < 18

- o Unable to provide informed consent

- o Pre-operative disorders in calcium homeostasis: hypo/hypercalcaemia with/without calcium supplementation

## Study design

### Design

**Study type:** Observational non invasive

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Diagnostic

### Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-02-2013

Enrollment: 220

Type: Anticipated

## Ethics review

Approved WMO

Date: 11-02-2014

Application type: First submission

Review commission: METC Erasmus MC, Universitair Medisch Centrum Rotterdam (Rotterdam)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

**In other registers**

Register	ID
CCMO	NL43050.078.12