

Critical investigation of the mechanisms at work in EMDR

Published: 17-08-2016

Last updated: 14-04-2024

The aim is to test whether EM in EMDR contributes to a reduction in dysfunctional beliefs about recalling the traumatic memory. A sample of Dutch veterans diagnosed with PTSD will be tested. During three EMDR treatment sessions we will measure...

Ethical review	Not approved
Status	Will not start
Health condition type	Anxiety disorders and symptoms
Study type	Interventional

Summary

ID

NL-OMON42938

Source

ToetsingOnline

Brief title

Working mechanisms in EMDR

Condition

- Anxiety disorders and symptoms

Synonym

Posttraumatic stress disorder; trauma

Research involving

Human

Sponsors and support

Primary sponsor: Universiteit Utrecht

Source(s) of monetary or material Support: Ministerie van Defensie

Intervention

Keyword: appraisal, EMDR, memory, trauma

Outcome measures

Primary outcome

Dysfunctional appraisals of memory recall.

Secondary outcome

Memory vividness and memory emotionality.

Study description

Background summary

Eye movement desensitization and reprocessing (EMDR) is a treatment-of-choice for posttraumatic stress disorder (PTSD) in many western countries. However, EMDR is not effective for about one-third of patients, and little is known about its underlying mechanisms. A unique component of EMDR therapy is that patients recall the traumatic memory and engage in horizontal eye movement (EM) simultaneously, which has been shown to add to its efficacy. How can this be explained? In recent years, growing attention has been paid to working memory theory. It holds that EM during recall of a (traumatic) memory causes a competition for limited working memory resources. By consequence, visual imagery becomes less vivid and less emotional. While this theory held up to rigorous testing, it is silent about how the immediate changes in memory phenomenology result in PTSD symptom reduction. Without such explanation, the clinical relevance of the existing knowledge is questionable. Two theoretical possibilities have been proposed. First, the direct loss in memory detail may be reconsolidated in long-term memory, causing future recollections to evoke less extreme emotional responses. Alternatively, it has been suggested that the empowering experience of holding a diluted traumatic memory in mind changes dysfunctional beliefs about the trauma memory (e.g., *I can now focus on it without becoming upset*), much like how exposure therapy works. The current research will test the latter hypothesis.

Study objective

The aim is to test whether EM in EMDR contributes to a reduction in dysfunctional beliefs about recalling the traumatic memory. A sample of Dutch veterans diagnosed with PTSD will be tested. During three EMDR treatment

sessions we will measure changes in memory vividness and emotionality (typical for experimental research on EMDR) and appraisals of memory recall (unique to the current research).

Study design

This study will cover the first three EMDR treatment sessions. A first session will provide baseline measurements. The second and third sessions will comprise recall of the trauma memory, one session with EM, the other one without. During each session participants will fill out a series of short questionnaires. By means of a cross-over experimental design for every participant one session (either session two or three, depending on randomised allocation) will be devoid of the eye movement component. As such, participants serve as their own control group.

Intervention

See "Study design": not applying the eye movements in one of the EMDR treatment sessions can be considered an intervention.

Study burden and risks

Burdens are that participants have to fill out questionnaires. Filling out the questionnaires may elicit feelings of discomfort as they refer to the trauma memory. There are no risks associated with participation.

Contacts

Public

Universiteit Utrecht

Heidelberglaan 1
Utrecht 3584CS
NL

Scientific

Universiteit Utrecht

Heidelberglaan 1
Utrecht 3584CS
NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years)

Elderly (65 years and older)

Inclusion criteria

Soldiers or veterans who developed PTSD, who are between 18-65 years old, and who have signed informed consent.

Exclusion criteria

High risk of suicide; psychotic disorders; dissociative experiences during course of the study.

Study design

Design

Study type:	Interventional
Intervention model:	Crossover
Masking:	Open (masking not used)
Control:	Uncontrolled
Primary purpose:	Treatment

Recruitment

NL	
Recruitment status:	Will not start
Enrollment:	25

Type: Anticipated

Ethics review

Not approved

Date: 17-08-2016

Application type: First submission

Review commission: METC Universitair Medisch Centrum Utrecht (Utrecht)

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register	ID
CCMO	NL58676.041.16