# Validation of adapted DCIEM Heliox diving tables using nitrox decompression

Published: 26-09-2014 Last updated: 20-04-2024

The proposed research intends to gather data to validate the hypothesis that using Nitrox to decompress during Heliox diving will significantly decrease the time and equipment requirements for decompression while being as safe as the standard heliox...

Ethical review	Approved WMO
Status	Recruitment stopped
Health condition type	Other condition
Study type	Observational non invasive

## Summary

## ID

NL-OMON47192

**Source** ToetsingOnline

Brief title validation of adapted DCIEM heliox tables

## Condition

• Other condition

**Synonym** decompression illness, decompression sickness

#### **Health condition**

hyperbare aandoeningen met name voorkoming decompressieziekten

#### **Research involving**

Human

## **Sponsors and support**

**Primary sponsor:** Koninklijke Marine **Source(s) of monetary or material Support:** Interspiro AB,Ministerie van Defensie;kosten METC toetsing wordt betaald door Industrie

### Intervention

Keyword: heliox diving, nitrox, tables, validation

## **Outcome measures**

#### **Primary outcome**

Changes in bubble scores as described by Kisman-Masurel (K-M) after the dive at

specified time points.

#### Secondary outcome

not applicable

# **Study description**

#### **Background summary**

For deep diving operations, Netherlands Navy divers use a breathing medium called Heliox, which can be breathed at a constant fraction of 16% oxygen and 84% helium. During decompression on Heliox there is a requirement to decompress on both air and oxygen. This type of diving is only done with Surface Supplied Diving systems where an umbilical (gas hose) delivers the breathing medium to the diver. The first stops during the decompression phase are conducted with air and once the diver reaches 9 metres of sea water (msw), 100% oxygen is used. This process therefore requires switching from the initial Heliox gas to air and then to oxygen to complete the decompression breathing gas could result in a saving of about 10% to 15% in overall decompression times compared with the standard heliox tables nowadays used. Furthermore only one switch in breathing gas is necessary. A feasibility study performed by DRDC seems to underline this hypothesis.

#### **Study objective**

The proposed research intends to gather data to validate the hypothesis that

2 - Validation of adapted DCIEM Heliox diving tables using nitrox decompression 22-05-2025

using Nitrox to decompress during Heliox diving will significantly decrease the time and equipment requirements for decompression while being as safe as the standard heliox tables nowadays used.

## Study design

Observational study

#### Study burden and risks

Burden: all divers have to make 3-6 dives. Before the dive they have to fill in a pre-dive questionnaire. During the dive they have to frequently score thermal comfort. After each dive they undergo Doppler measurements for at least 3 times.

Benefits: For military diving, it is of importance to use safe diving tables. Furthermore, diving tables with shorter decompression time could are operational beneficial. However these shorter decompression times should not lead to a higher incidence of decompression sickness (DCS).

Risks assessment, group relatedness: The risk of DCS associated with this dive simulation can be considered as very small (< 1.0%). When a diver encounters symptoms of DCS he will be treated according to the guidelines used within the Royal Netherlands Navy, which are equal to those internationally used. Beside the risk of decompression sickness, there is the risk for central nervous system oxygen toxicity, which is estimated as < 0.01%. Furthermore, as described in the information package subjects participating in this trial run a small risk on hypothermia, hypercapnia, and hypoxia. All in all, we think the risk of these simulations can be considered low, and equal to a real, wet operational heliox dive.

# Contacts

**Public** Koninklijke Marine

Rijkszee- en Marinehaven 1 Den Helder 1781 zz NL **Scientific** Koninklijke Marine

Rijkszee- en Marinehaven 1 Den Helder 1781 zz NL

3 - Validation of adapted DCIEM Heliox diving tables using nitrox decompression 22-05-2025

# **Trial sites**

## **Listed location countries**

Netherlands

# **Eligibility criteria**

Age Adults (18-64 years) Elderly (65 years and older)

## **Inclusion criteria**

adult males, professional divers, medically fit to dive

## **Exclusion criteria**

if one of the inclusion criteria is not met, recent infection and/or flue, any previous decompression sickness, daily use of alcoholic beverage

# Study design

## Design

Study type: Observational non invasive		
Masking:	Open (masking not used)	
Control:	Uncontrolled	
Primary purpose:	Prevention	

## Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	17-02-2015
Enrollment:	50

4 - Validation of adapted DCIEM Heliox diving tables using nitrox decompression 22-05-2025

#### Actual

# **Ethics review**

Approved WMO Date:	26-09-2014
Application type:	First submission
Review commission:	METC Amsterdam UMC
Approved WMO Date:	04-04-2018
Application type:	Amendment
Review commission:	METC Amsterdam UMC

# **Study registrations**

## Followed up by the following (possibly more current) registration

No registrations found.

## Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

 Register
 ID

 CCMO
 NL48626.018.14

# **Study results**

Date completed:	12-10-2018
Actual enrolment:	53