

# A 12-week randomized, controlled trial to compare TOUJEO® and TRESIBA® in terms of glucose values in target range and variability during continuous glucose monitoring in patients with type 1 diabetes mellitus

Published: 16-10-2019

Last updated: 10-04-2024

To demonstrate the noninferiority of insulin glargine 300 U/mL in comparison to insulin degludec 100 U/mL on glycemic control and variability in participants with diabetes mellitus

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Diabetic complications
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON48426

### Source

ToetsingOnline

### Brief title

inRange

### Condition

- Diabetic complications

### Synonym

Diabetes Type I, glucose values

### Research involving

Human

## Sponsors and support

**Primary sponsor:** Sanofi-aventis

**Source(s) of monetary or material Support:** Sanofi

## Intervention

**Keyword:** diabetes type I, Glucose monitoring, LPS14947

## Outcome measures

### Primary outcome

Percent time in glucose range of \*70 to \*180 mg/dL (\*3.9 to \*10 mmol/L) at Week 12, obtained using continuous glucose monitoring (CGM)

### Secondary outcome

At Week 12, the following endpoints will be assessed:

- \* Glucose total CV
- \* Glucose within-day and between-day CV

Other secondary endpoints:

- \* Change from baseline to Week 12 in HbA1c
- \* Change from baseline to Week 12 in central lab FPG
- \* Percent time and mean hours per day with glucose<70mg/dL (on all time and during the night [00:00 to 05:59])
- \* Percent time and mean hours per day with glucose>180mg/dL

Number of participants with adverse events (see Section 8.3)

Number of participants with at least one hypoglycemic event

from baseline to Week 12

Number of hypoglycemic events per participant year from

baseline to Week 12

## Study description

### Background summary

The fluctuations in blood glucose, regular short-duration periods of hyperglycemia and hypoglycemia, which are not detected by the measurements of glycated hemoglobin (HbA1c), may possibly contribute to vascular pathological processes and diabetic complications.

Insulin glargine U300 (HOE901-U300, Toujeo®) is a more concentrated formulation of insulin

glargine U100 (HOE901) which showed a prolonged and flatter glucose-lowering activity (up to

36 hours) compared to Lantus (Insulin glargine U100), resulting in less circadian fluctuation in

blood glucose levels (1). Given these features of the pharmacokinetic (PK)/pharmacodynamics

(PD) profiles, Toujeo is considered to be well suitable for constant, peakless 24-hour basal insulin

supply in diabetes management, with the expectation of less hypoglycemic risk at equal to tighter

blood glucose control.

### Study objective

To demonstrate the noninferiority of insulin glargine 300 U/mL in comparison to insulin degludec 100 U/mL on glycemic control and variability in participants with diabetes mellitus

### Study design

This is a multicenter, randomized, active-controlled, parallel-group, 12-week open-label study to

compare the efficacy of Toujeo with Tresiba using 20-day CGM glucose profiles at Week 12. The

CGM during study period will be blinded to both the Investigators and patients

## Intervention

Patients taking their current basal insulin at any time (eg, evening) other than in the morning should switch their injection time to the morning at randomization.

## Study burden and risks

because participants are frequently monitored for blood glucose values, the risk of hyperglycemia and hypoglykemie are low.  
Participants should be regularly monitored for 12 weeks and need 2x 20 days to wear a continuous glucose meter with them. minimum blood samples can be considered standard of care for this group of participants

## Contacts

### Public

Sanofi-aventis

Paasheuvelweg 25  
Amsterdam 1105 BP  
NL

### Scientific

Sanofi-aventis

Paasheuvelweg 25  
Amsterdam 1105 BP  
NL

## Trial sites

### Listed location countries

Netherlands

## Eligibility criteria

### Age

Adults (18-64 years)

Elderly (65 years and older)

## Inclusion criteria

Participants with Type 1 Diabetes mellitus (T1DM)

Participants treated with multiple daily injections (MDI) using basal insulin analog once daily and rapid acting insulin analogs for at least one year

HbA1c \* 7% (48 mmol/mol) and \* 10% (86 mmol/mol) at screening

## Exclusion criteria

Participants not on stable dose of basal insulin analog

Participants having received Toujeo or Tresiba as basal insulin within 30 days prior to screening

Participants not using the same insulins (both basal and rapid) within 30 days prior to screening

Participants having received basal insulin dose \*0.6 U/kg body weight within 30 days prior to screening

Participants having received any glucose lowering drugs (including any premixed insulins, human regular insulin as mealtime insulins, any others injectable or oral), other than basal and rapid insulin analogs, within 3 months prior to screening

End stage renal disease or on renal replacement treatment

Retinopathy or maculopathy with one of the following treatments, either recent (within 3 months prior to screening) or planned: intravitreal injections or laser or vitrectomy surgery

Body weight change \*5 kg within 3 months prior to screening

## Study design

### Design

Study phase: 4

Study type: Interventional

Intervention model:	Parallel
Allocation:	Randomized controlled trial
Masking:	Open (masking not used)
Control:	Active
Primary purpose:	Treatment

## Recruitment

NL	
Recruitment status:	Recruitment stopped
Start date (anticipated):	25-11-2020
Enrollment:	45
Type:	Actual

## Medical products/devices used

Product type:	Medicine
Brand name:	Toujeo
Generic name:	insulin glargine
Registration:	Yes - NL intended use
Product type:	Medicine
Brand name:	Tresiba
Generic name:	insulin degludec
Registration:	Yes - NL intended use

## Ethics review

Approved WMO	
Date:	16-10-2019
Application type:	First submission
Review commission:	BEBO: Stichting Beoordeling Ethiek Bio-Medisch Onderzoek (Assen)
Approved WMO	
Date:	11-12-2019
Application type:	First submission
Review commission:	BEBO: Stichting Beoordeling Ethiek Bio-Medisch Onderzoek (Assen)

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
EudraCT	EUCTR2017-002756-91-NL
CCMO	NL71287.056.19