

# The Swallowing Exercise Aid in Total Laryngectomy: Feasibility, Adherence and a Multidimensional Assessment of Functional-, Physical-, and Patient-reported outcomes

Published: 27-10-2021

Last updated: 12-10-2024

The primary objective is to evaluate the feasibility and adherence of 6 weeks of rehabilitation with the SEA in participants who underwent TL. The secondary objective is to examine if the SEA is an effective tool to improve swallowing function and...

<b>Ethical review</b>	Approved WMO
<b>Status</b>	Recruitment stopped
<b>Health condition type</b>	Other condition
<b>Study type</b>	Interventional

## Summary

### ID

NL-OMON50048

### Source

ToetsingOnline

### Brief title

SEATTLE

### Condition

- Other condition

### Synonym

dysphagia, Larynx extirpation

### Health condition

Hoofdhalskankerpatienten die een totale laryngectomie hebben ondergaan

## Research involving

Human

## Sponsors and support

**Primary sponsor:** Antoni van Leeuwenhoek Ziekenhuis

**Source(s) of monetary or material Support:** Unrestricted research grant Atos Medical AB  
Zweden

## Intervention

**Keyword:** MRI, Swallowing training, Total laryngectomy

## Outcome measures

### Primary outcome

This study has a multidimensional assessment approach, including feasibility and adherence, physical outcomes, functional outcomes, and patient-reported outcomes. In this study, measures of interest are muscle strength, manofluorography, manometry, voice recordings, dynamic range of voice, maximum fonation time, questionnaires, and information from the electronic medical files.

### Secondary outcome

Secondary: (Multidimensional Evaluation)

Physical outcomes:

- Strength measurement (IOPI, dynamometer)
- Manofluorography
- Pressure measurement during breathing and vocalization (manometry)

Functional Outcomes:

- Voice recordings
- Mouth opening

Subjective outcomes:

- Questionnaires on Quality of life, swallowing problems, voice problems, participation

## Study description

### Background summary

Total laryngectomy (TL), implicating surgical removal of the larynx, is the operation of choice in patients with advanced laryngeal or hypopharyngeal cancer. Functional disabilities associated with TL include impairment of swallowing, speech, olfaction, and taste. Muscle strength, muscle coordination, and pressure build-up are essential factors for swallowing and tracheoesophageal speech. Research has shown that TL negatively influences the patients' pressure build-up at the base of the tongue and coordination of muscular contraction in the neopharynx. The Swallow Exercise Aid (SEA) is developed as an exercise tool that activates and strengthens the suprahyoid and pharyngeal muscles relevant for swallowing function. The feasibility and adherence were considered good in healthy senior adults and head and neck cancer patients with chronic dysphagia. Following these positive results, the feasibility, adherence, and effectiveness of its use for rehabilitation of swallowing and tracheoesophageal speech will be studied in patients with altered anatomy because of TL.

### Study objective

The primary objective is to evaluate the feasibility and adherence of 6 weeks of rehabilitation with the SEA in participants who underwent TL. The secondary objective is to examine if the SEA is an effective tool to improve swallowing function and tracheoesophageal speech. The outcomes include functional (swallowing efficiency, vocal functioning) and physical (intra-pharyngeal pressure) outcomes, patient-reported complaints, and quality of life.

### Study design

The proposed study is a pilot study and includes a prospective feasibility study involving one training group, six weeks of training and three moments of assessment.

### Intervention

All participants have to train three different strengthening exercises with the

SEA three times a day, including the Chin Tuck against resistance (CTAR+), Jaw Opening against resistance (JOAR+), and the Effortful swallow exercise (ES+) for six weeks.

### **Study burden and risks**

Participating in this study and training with the SEA might improve swallowing function and voice quality. These improvements may lead to enhance the quality of life. Muscular pain or muscular exhaustion might be the only risk associated with participation in this study. The assessments contain radiation and manometry in which nose bleeding is a minimal risk.

## **Contacts**

### **Public**

Antoni van Leeuwenhoek Ziekenhuis

Plesmanlaan 121  
Amsterdam 1066CX  
NL

### **Scientific**

Antoni van Leeuwenhoek Ziekenhuis

Plesmanlaan 121  
Amsterdam 1066CX  
NL

## **Trial sites**

### **Listed location countries**

Netherlands

## **Eligibility criteria**

### **Age**

Adults (18-64 years)

Elderly (65 years and older)

## Inclusion criteria

- $\geq 18$  years or older
- Undergone a TLE
- At least six months post TLE
- At least six months postoperative (chemo-) radiotherapy (if applicable)
- Using Tracheoesophageal speech
- Oral intake of a modified diet (FOIS 2 - 6) or who experience swallowing problems
- Signed informed consent

## Exclusion criteria

- Unable to understand the patient information
- Unable to understand the informed consent
- Unable to comprehend and use the SEA
- Unable or unwilling to provide informed consent
- Neuro-degenerative diseases, for instance, dementia, Parkinson, Korsakov

## Study design

### Design

**Study type:** Interventional

Masking: Open (masking not used)

Control: Uncontrolled

Primary purpose: Treatment

### Recruitment

NL

Recruitment status: Recruitment stopped

Start date (anticipated): 21-04-2022

Enrollment: 20

Type: Actual

## Medical products/devices used

Generic name: Swallowing Exercise Aid (SEA)

Registration: Yes - CE intended use

## Ethics review

Approved WMO  
Date: 27-10-2021  
Application type: First submission  
Review commission: METC NedMec

## Study registrations

### Followed up by the following (possibly more current) registration

No registrations found.

### Other (possibly less up-to-date) registrations in this register

No registrations found.

### In other registers

Register	ID
CCMO	NL78528.031.21

## Study results

Date completed: 16-10-2023