Posterior Tibial Tendon Transfer in Foot Drop; Description and Analysis of the Circumtibial Route

Published: 26-03-2024 Last updated: 04-06-2024

To assess the range and strength of dorsiflexion of the ankle and the percentage of AFO-use after executing a variation on the circumtibial route of the posterior tibial tendon transfer, performed in our center on patients with foot drop.

Ethical review	Not approved
Status	Will not start
Health condition type	Peripheral neuropathies
Study type	Observational non invasive

Summary

ID

NL-OMON56697

Source ToetsingOnline

Brief title Posterior Tibial Tendon Transfer in Foot Drop

Condition

- Peripheral neuropathies
- Bone and joint therapeutic procedures

Synonym Foot drop, peroneal nerve palsy

Research involving Human

Sponsors and support

Primary sponsor: Leids Universitair Medisch Centrum Source(s) of monetary or material Support: Ministerie van OC&W

1 - Posterior Tibial Tendon Transfer in Foot Drop; Description and Analysis of the ... 14-06-2025

Intervention

Keyword: Foot drop, Orthopedic surgery, Peroneal palsy, Tendon transfer

Outcome measures

Primary outcome

The need for AFO after surgery compared to before surgery.

Secondary outcome

Active dorsiflexion of the ankle after tendon transfer surgery compared to

before surgery. Furthermore, pain, range of motion of the ankle, muscle

strength of the ankle, limitations during daily life activities and

complications

Study description

Background summary

Foot drop is an invalidating loss of dorsiflexion of the ankle as a result of damage of the peroneal nerve. For years a transfer of the posterior tibial tendon has been a surgical options for patients with foot drop. The goal of this type of surgery is to restore dorsiflexion limiting the dependence from ankle-foot orthoses (AFO). Most surgeons perform a tibial tendon transfer trough the interosseous membrane. As an alternative, the circumtibial route of posterior tibial tendon transfers has been described in patients suffering from leprosy. This technique is easier to perform, and has potentially similar results with a lower complication rate. In our center, the circumtibial route has been the standard surgical approach, but literature lacks studies in traumatic foot drop patients. Our hypothesis is that the circumtibial route offers a safe surgical technique that significantly restores ankle dorsiflexion resulting in a reduced dependency on AFO.

Study objective

To assess the range and strength of dorsiflexion of the ankle and the percentage of AFO-use after executing a variation on the circumtibial route of the posterior tibial tendon transfer, performed in our center on patients with

foot drop.

Study design

Case series

Study burden and risks

There are no risks associated with participation.

Contacts

Public Leids Universitair Medisch Centrum

Albinusdreef 2 Leiden 2333 ZA NL **Scientific** Leids Universitair Medisch Centrum

Albinusdreef 2 Leiden 2333 ZA NL

Trial sites

Listed location countries

Netherlands

Eligibility criteria

Age

Adults (18-64 years) Elderly (65 years and older)

Inclusion criteria

3 - Posterior Tibial Tendon Transfer in Foot Drop; Description and Analysis of the ... 14-06-2025

Patient with a drop foot due to a traumatic peripheral nerve injury treated with a posterior tibial tendon transfer.

Exclusion criteria

- No informed consent
- Central neurological disorders (e.g. cerebrovascular events, leprosy)
- Previous surgery such as ankle arthrodeses or other operations or disabilities influencing ankle function.
- History of a malignancy in the affected limb
- History of muscle dystrophy or metabolic disease
- History of rheumatoid arthritis
- Insufficient Dutch language skills.

Study design

Design

Study type: Observational non invasive		
Masking:	Open (masking not used)	
Control:	Uncontrolled	
Primary purpose:	Treatment	

Recruitment

NL	
Recruitment status:	Will not start
Enrollment:	30
Туре:	Anticipated

Ethics review

Not approved	
Date:	26-03-2024
Application type:	First submission
Review commission:	METC Leiden-Den Haag-Delft (Leiden)
	metc-ldd@lumc.nl

4 - Posterior Tibial Tendon Transfer in Foot Drop; Description and Analysis of the ... 14-06-2025

Study registrations

Followed up by the following (possibly more current) registration

No registrations found.

Other (possibly less up-to-date) registrations in this register

No registrations found.

In other registers

Register CCMO **ID** NL84161.058.23