# Effect of Plant Based High Energy High Protein Oral Nutritional Supplement on Nutritional Intake in Patients with or at Risk of Disease Related Malnutrition.

Published: 04-12-2024 Last updated: 18-01-2025

To investigate the effect of PB HEHP ONS compared to DB HEHP ONS on nutritional intake (energy & protein) in subjects with or at risk of DRM and/or prescribed with ONS.

Ethical review Approved WMO

**Status** Pending

**Health condition type** Other condition **Study type** Interventional

## **Summary**

#### ID

NL-OMON57157

**Source** 

ToetsingOnline

**Brief title** 

**PRO-PLANT** 

#### **Condition**

Other condition

#### **Synonym**

disease related malnutrition / malnutrition

#### **Health condition**

Ziekte gerelateerde ondervoeding

#### Research involving

Human

### **Sponsors and support**

Primary sponsor: Danone Global R&I Center

**Source(s) of monetary or material Support:** Danone Global Research & Innovation

Center B.V.

#### Intervention

Keyword: Disease Related Malnutrition (DRM), Oral nutritional support (ONS), Plant-Based

#### **Outcome measures**

#### **Primary outcome**

Main outcome of the study:

To investigate the effect of PB HEHP ONS compared to DB HEHP ONS on nutritional intake (energy & protein) in subjects with or at risk of Disease Related Malnutrition (DRM) and/or prescribed with ONS.\*

#### **Secondary outcome**

other outcomes of the study:

To investigate the effect of PB HEHP ONS compared to DB HEHP ONS in subjects with or at risk of DRM and/or prescribed with ONS prior to the study on:\*

- Safety and gastrointestinal (GI) tolerance\*
- Compliance\*\*
- Product experience & liking\*
- Changes in anthropometry (BW, BMI)\*
  - 2 Effect of Plant Based High Energy High Protein Oral Nutritional Supplement on Nu ... 25-06-2025

- Changes in blood nutrient status (Vitamin D (25(OH)D), ferritin, zinc, calcium, parathyroid hormone (PTH)\*
- Changes in percentage of subjects with or at risk of malnutrition according to the Malnutrition Universal Screening Tool (MUST).\*
- Changes in health-related quality of life (QoL) (EQ-5D-5L)\*
- Changes in muscle strength & function (Hand Grip Strength & x 5 sit to stand test).\*\*

# **Study description**

#### **Background summary**

The World Health Organization (WHO) reports the global prevalence of malnutrition showing that 462 million adults and about 2 billion people are underweight and suffer from micronutrient deficiency, respectively. According to the Medical Nutrition International Industry dossier the estimated number of people suffering from malnutrition in Europe is 33 million (MNI 2018). The prevalence of malnutrition can be associated with specific groups of disease and settings of care. Disease related malnutrition (DRM) is the consequence of inadequate dietary intake and nutritional support, and it can negatively impact the recovery of patients. Oral nutritional supplements (ONS) have proven clinical benefits for malnourished patients: ONS use is linked to lower mortality and lower complication rates when compared to standard of care. Historically, most ONS are made with dairy proteins, however, as more people transition to PB diets and the demand for products that do not contain animal derived ingredients grows, there is a growing request for PB ONS to meet this unmet need. Currently there are some PB ONS on the market, but very few that are high in energy and high in protein, and ready to drink. There is also very little evidence currently on plant based medical nutrition.\*

#### **Study objective**

To investigate the effect of PB HEHP ONS compared to DB HEHP ONS on nutritional intake (energy & protein) in subjects with or at risk of DRM and/or prescribed with ONS.

#### Study design

This is a randomised controlled, double blind, parallel-group, multi-centre, multi-country, exploratory study.

#### Intervention

Subjects that are eligible for study participation will be randomly allocated to receive either the Test Product or the Control Product for a period of 8 weeks.

#### Study burden and risks

The burden on the subjects is kept minimal. Subjects are mainly asked to complete questionnaires and diaries during screening and/or the intervention period. The expected risks associated with the test and control products are minimal. During the intervention period, the number of visits to the research center (3x) and phone calls (2x) is limited. In consultation and if the site can facilitate (i.e. mobile blood collection), the visits can also take place at the participants' homes in order to relieve the participants. A large patient population can benefit from the results of the research.

## **Contacts**

#### **Public**

Danone Global R&I Center

Uppsalalaan 12 Utrecht 3584 CT NL

#### Scientific

Danone Global R&I Center

Uppsalalaan 12 Utrecht 3584 CT NL

## **Trial sites**

#### **Listed location countries**

**Netherlands** 

# **Eligibility criteria**

#### Age

Adults (18-64 years)

#### Inclusion criteria

- 1. Age >= 18 years
- 2.Identified as at medium or high risk of malnutrition based on:
- a) MUST score >= 1 and / or b)Being prescribed with ONS and willing and able to switch from pre-study prescribed ONS to the Test or Control Product for participation in the study
- 3. In need of 2 servings of ONS/day (400 kcal; 20 gr protein per serving) for at least 8 weeks as determined by the treating healthcare professional.
- 4. Medically and physically able to consume high energy high protein ONS in the opinion of the Investigator.
- 5. Willing to maintain dietary habits for the duration of the study.
- 6. Willing to consume plant based as well as dairy based ONS.

#### **Exclusion criteria**

- 1. Known allergy to soy, cow\*s milk protein or to any other ingredients as listed in the study product composition (refer to the product information brochure (PIB) and the appendix of this protocol).
- 2. Known intolerance to any ingredients as listed in the study product composition (refer to PIB and appendix of this protocol). Subjects with lactose intolerance who use lactase may still be enrolled in the study.
- 3. Known galactosaemia
- 4. Any contraindication to oral feeding per se being: gastrointestinal failure or suppressed gastrointestinal function, complete intestinal obstruction and major intra-abdominal sepsis.
- 5. Active flare of inflammatory bowel disease as defined by Harvey-Bradshaw Index (HBI) >6 (Crohn\*s disease) or Simple Clinical Colitis Activity Index (SCCAI) >5 (ulcerative colitis).
- 6. Recent history of gastrointestinal surgery (except appendectomy) that

interferes with the GI function, e.g. ileostomy, colostomy, (partial) gastrectomy or any other procedure for stomach volume reduction including gastric banding, within four weeks prior to the screening visit.

- 7. Requiring a protein restricted diet as confirmed by a physician, for example chronic kidney disease stage 4 and 5 (estimated by Glomerular Filtration Rate <30 mL/min/1.73 m2).
- 8. End stage renal disease (with renal replacement therapy e.g. haemodialysis or peritoneal dialysis), who may not require protein restriction.
- 9. Requiring enteral nutrition (via tube delivery) or parenteral nutrition.

# Study design

## **Design**

Study type: Interventional

Intervention model: Parallel

Allocation: Randomized controlled trial

Masking: Double blinded (masking used)

Control: Active

Primary purpose: Other

#### Recruitment

NL

Recruitment status: Pending

Start date (anticipated): 01-02-2025

Enrollment: 40

Type: Anticipated

## Medical products/devices used

Registration: No

## **Ethics review**

Approved WMO

Date: 04-12-2024

Application type: First submission

Review commission: BEBO: Stichting Beoordeling Ethiek Bio-Medisch Onderzoek

(Assen)

# **Study registrations**

# Followed up by the following (possibly more current) registration

No registrations found.

## Other (possibly less up-to-date) registrations in this register

No registrations found.

## In other registers

Register ID

CCMO NL87757.056.24

Other Registratie in Clintrials.gov database volgt