

# Enterovirus- and parechovirus infection in children.

Gepubliceerd: 12-12-2011 Laatste bijgewerkt: 15-05-2024

To determine the epidemiology, diagnosis and prognosis of enterovirus (EV) and parechovirus (HPeV) infection in Dutch children. Hypotheses: 1. EVs are one of the most important viral agents of infection in Dutch children; 2. EVs are the major...

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving gestart
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Observationeel onderzoek, zonder invasieve metingen

## Samenvatting

### ID

NL-OMON20807

### Bron

Nationaal Trial Register

### Aandoening

Enterovirus (EV), a picornavirus, is a common cause of infection in children. The incidence of EV infection in Dutch children is not exactly known. There is no official registration. EVs cause a broad range of clinical syndromes from gastro-enteritis to meningitis. The clinical presentation changes between the several subtypes. Human Parechovirus (HPeV), also a member of the Picornaviridae, is recently identified, and is associated with similar symptoms as EV infection. EV and HPeV can be diagnosed with viral cultures, reverse transcription polymerase chain reaction (PCR) of feces, urine, throat swab, blood or cerebrospinal fluid (CSF) or with serology. Not much is known about the prognosis of enterovirus and parechovirus menigitis in children.

### Ondersteuning

**Primaire sponsor:** This study was started in the St. Elisabeth Hospital Tilburg, the Netherlands, Pediatrics, C. Obihara.

There are no sponsors or fundings.

**Overige ondersteuning:** No fundings.

# Onderzoeksproduct en/of interventie

## Uitkomstmaten

### Primaire uitkomstmaten

1. Specific clinical symptoms in patients with proven EV or HPeV infection;<br>
2. Different laboratory diagnostic methods: viral culture, PCR and serology;<br>
3. Several body fluids: feces, urine, throat swab, blood, and eventually CSF;<br>
4. Developmental milestones of children after an EV/HPeV CNS infection.

## Toelichting onderzoek

### Achtergrond van het onderzoek

Patients are included only in the Netherlands (St. Elisabeth hospital Tilburg, Tweesteden hospital Tilburg, Amphia Hospital Breda).

Background of the study:

Enterovirus (EV), a picornavirus, is a common cause of infection in children. The incidence of EV infection in Dutch children is not exactly known. There is no official registration. EVs cause a broad range of clinical syndromes from gastro-enteritis to meningitis. The clinical presentation changes between the several subtypes. Human Parechovirus (HPeV), also a member of the Picornaviridae, is recently identified, and is associated with similar symptoms as EV infection. EV and HPeV can be diagnosed with viral cultures, reverse transcription polymerase chain reaction (PCR) of feces, urine, throat swab, blood or cerebrospinal fluid (CSF) or with serology.

Hypotheses:

- (1) EVs are one of the most important viral agents of infection in Dutch children;
- (2) EVs are the major cause of meningitis in Dutch children;
- (3) PCR is more sensitive than viral culture and serology in the detection of EV and HPeV infection in children;
- (4) HPeV is a major cause of serious infection in younger children ( $\leq 2$  year) with

psychomotoric and cognitive develop deficiencies.

Objective of the study:

Primary objectives:

- (1) To describe the incidence of EV and HPeV infections in Dutch children;
- (2) To determine the major symptoms of EV/HPeV infections in children;
- (3) To evaluate the epidemiology and symptoms of the different subtypes of EV and HPeV;
- (4) To compare the sensitivity and specificity of different laboratory techniques to detect EV or HPeV during infection: viral culture, PCR and serology;
- (5) To determine the sensitivity and specificity of the different body fluids in diagnosis of EV or HPeV infection: feces, urine, throat swab, blood, CSF;
- (6) To determine the sequelae after an EV/HPeV central nervous system (CNS) infection till 5 years after infection.

Secondary objectives:

- (1) To describe the morbidity of EV and HPeV infection (hospitalization, duration of hospitalization, duration and severity of symptoms, average school- and work absence, duration use of antibiotics);
- (2) To evaluate the use of antibiotics before and after the diagnosis.

Study design:

Observational multicenter study with nested case-control follow-up study. To identify EV or HPeV infection we will examine feces, urine, throat swab, blood and CSF.

The definition of cases and controle in the follow-up study is as follows:

Cases: Children with an CNS infection with EN/HPeV, proven with EV/HPeV in liquor (meningitis or encephalitis).

Control group 1: No EV/HPeV meningitis, however EV/HPeV positivity in feces, urine, throat swab or blood.

Control group 2: No EV/HPeV or other viral (or bacterial) infection proven Approximately 2 weeks after the infection, the patient will be invited for check-up for a standardized questionnaire, physical examination and for the second vena puncture. The further follow-up will consist of taking standardized questionnaires and developmental and cognitive tests at 6, 12, 24 and 60 months after the infection.

Study population:

Children  $\leq$  16 years of age with suspected EV or HPeV infection.

Primary study parameters/outcome of the study:

1. Specific clinical symptoms in patients with proven EV or HPeV infection;
2. Different laboratory diagnostic methods: viral culture, PCR and serology;
3. Several body fluids: feces, urine, throat swab, blood, and eventually CSF;
4. Developmental milestones of children after an EV/HPeV CNS infection.

Secondary study parameters/outcome of the study (if applicable):

1. Morbidity due to EV/HPeV infection (hospitalization, duration of hospitalization, duration and severity of symptoms, average school- and work absence, duration use of antibiotics);
2. Antibiotic treatment of patients (antibiotics, stop of antibiotics after diagnosis of EV or HPeV infection).

## **Doel van het onderzoek**

To determine the epidemiology, diagnosis and prognosis of enterovirus (EV) and parechovirus (HPeV) infection in Dutch children.

Hypotheses:

1. EVs are one of the most important viral agents of infection in Dutch children;
2. EVs are the major cause of meningitis in Dutch children;

3. PCR is more sensitive than viral culture and serology in the detection of EV and HPeV infection in children;

4. HPeV is a major cause of serious infection in younger children ( $\leq 2$  year) with psychomotoric and cognitive develop deficiencies.

### **Onderzoeksopzet**

4 follow up visits at 6, 12, 24 and 60 months after infection for standardized questionnaire, physical examination and developmental tests.

Psychotherapist: m-ABC, BSID-II.

Psychologist: BSID, CBCL, RAKIT/WPSII.

### **Onderzoeksproduct en/of interventie**

When children are participating in the study the following materials are collected:

1. Feces;
2. Urine;
3. Nasopharynx swab;
4. Blood sample (if parents agree);
5. Cerebrospinal fluid, ONLY if the pediatrician has a suspicion of a meningitis and the pediatrician says a spinal puncture is indicated.

For the follow up patients are classified in one of the following groups:

1. Group 1: Children with an EV or HPeV meningitis;
2. Group 2: Children with an EV or HPeV infection but no meningitis;
3. Group 3: Children suspected for an EV or HPeV infection, but there was no proven cause of infection (no EV, HPeV or other virus/bacteria was found).

For the follow up patients are seen in our outpatient clinic 2 weeks after the infection by the pediatrician. Furthermore there are 4 follow up visits at 6, 12, 24 and 60 months after infection for standardized questionnaire, physical examination and developmental tests (m-ABC/ BSID-II). They are seen by the pediatrician, psychoterapist and psychologist.

## Contactpersonen

### Publiek

St. Elisabeth Ziekenhuis Tilburg<br>  
Afdeling kindergeneeskunde<br>  
Hilvarenbeekseweg 60  
Stephanie Crom, de  
Tilburg 5022 GC  
The Netherlands

### Wetenschappelijk

St. Elisabeth Ziekenhuis Tilburg<br>  
Afdeling kindergeneeskunde<br>  
Hilvarenbeekseweg 60  
Stephanie Crom, de  
Tilburg 5022 GC  
The Netherlands

## Deelname eisen

### Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

A + B are required.

A. All Children  $\leq$  16 years of age with a clinical suspicion of an EV/HPeV infection:

1. Fever (temperature  $\geq 38,0$  degrees C ) OR;
2. Meningeal inflammation: (Anamnestic or with examination at least 2 of the following: headache, photophobia, nuchal rigidity, irritability, lethargy, nausea, vomiting, drowsiness, positive sign of Kernig or Brudzinsky) OR;
3. At least 3 of the following: Hypothermia, headache, drowsiness, nuchal rigidity, irritability,

photofobia, vomiting, diarrhea, anorexia, coughing, myalgia, rash OR;

4. Sepsis: Clinical suspicion of an infection plus: - temperature  $>38,5$  degrees C of  $< 36$  degrees C rectal or oral - tachycardia: heart rate  $> 2$  SD for age OR children  $< 1$  year with a bradycardia: heart rate  $< 2$  SD for age - tachypnea: breathing rate  $> 2$  SD for age.

B. Signed informed consent by the parent(s)/legal guardian(s).

## **Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)**

1. Other proven cause of the infection: Positive bacterial, viral (other than EV/HPeV), parasitic or fungal/yeast culture or PCR (feces, urine, throat swab, blood, CSF);
2. Other causes of illness: Neoplasma, auto-immune diseases, rheumatic diseases, endocrinologic diseases, gastroesophageal reflux, etcetera;
3. Known psycho-motor retardation, metabolic diseases with neuro-muscular or cognitive abnormalities;
4. Patients older than 16 years of age;
5. No signed informed consent from the parent(s)/ legal Guardian(s).

## **Onderzoeksopzet**

### **Opzet**

Type:	Observationeel onderzoek, zonder invasieve metingen
Onderzoeksmodel:	Factorieel
Toewijzing:	N.v.t. / één studie arm
<b>Controle:</b>	N.v.t. / onbekend

### **Deelname**

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	17-03-2008
Aantal proefpersonen:	240
Type:	Verwachte startdatum

## Ethische beoordeling

Positief advies

Datum: 12-12-2011

Soort: Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 31728

Bron: ToetsingOnline

Titel:

### Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

### In overige registers

Register	ID
NTR-new	NL3045
NTR-old	NTR3193
CCMO	NL21361.008.07
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON31728

## Resultaten

### Samenvatting resultaten

N/A