

# Can NT-proBNP guided therapy during hospital admission for acutely decompensated heart failure reduce mortality and readmissions?

Gepubliceerd: 08-02-2012 Laatste bijgewerkt: 19-03-2025

NT-proBNP guidance during in-hospital treatment of acutely decompensated heart failure (to strive for >30% reduction) reduces readmissions and mortality and increases the number of days alive out of hospital in the first 180 days.

<b>Ethische beoordeling</b>	Positief advies
<b>Status</b>	Werving gestopt
<b>Type aandoening</b>	-
<b>Onderzoekstype</b>	Interventie onderzoek

## Samenvatting

### ID

NL-OMON21346

### Bron

Nationaal Trial Register

### Verkorte titel

PRIMA II study

### Aandoening

Acutely decompensated heart failure

Acute heart failure

Acuut hartfalen

Acuut gedecompenseerd hartfalen

### Ondersteuning

**Primaire sponsor:** Academical Medical Center - University of Amsterdam

**Overige ondersteuning:** The Dutch Heart Foundation

Roche Diagnostics BV

# Onderzoeksproduct en/of interventie

## Uitkomstmaten

### Primaire uitkomstmaten

1. Readmissions and mortality in the first 180 days;<br>
2. The number of days alive out of hospital in the first 180 days.

## Toelichting onderzoek

### Achtergrond van het onderzoek

Guiding therapy of heart failure (HF) by an objective measure like NT-proBNP has received intense attention. The gain that is made by this form of guidance is modest when applied to chronic heart failure (CHF) patients. Recent post discharge data from our own group does show that NT-proBNP guidance can detect important short-term changes. Studies have also shown that NT-proBNP discharge value and a >30% NT-proBNP reduction during admission are statistically significant predictors of readmissions and mortality. These data suggest a role for such NT-proBNP guidance, rather in an acute than in a chronic setting. Acute admission for HF occurs frequent: in 2004, there were almost 25,000 hospital admissions in the Netherlands. Particularly worrisome is the high percentage of readmissions which reaches 30 to 60% within 6 months, importantly increasing the economic burden of this disease. In short, in-hospital care for acutely decompensated heart failure may be improved by NT-proBNP guidance to reduce the number of readmissions.

The primary objective of the present multicenter randomized controlled trial is to demonstrate that NT-proBNP guidance during in-hospital treatment for acutely decompensated heart failure (to strive for >30% reduction) reduces readmissions and mortality and increases the number of days alive out of hospital in the first 180 days compared to therapy guided by standard clinical judgment. Morbidity and mortality is measured in terms of days alive outside the hospital within the follow-up period of 180 days.

### Doel van het onderzoek

NT-proBNP guidance during in-hospital treatment of acutely decompensated heart failure (to strive for >30% reduction) reduces readmissions and mortality and increases the number of days alive out of hospital in the first 180 days.

### Onderzoeksopzet

Endpoint assessment (readmission or death) will take place at 4 timepoints after discharge: 1 week, 1 month, 3 months and 6 months after discharge.

### **Onderzoeksproduct en/of interventie**

1. NT-proBNP guided therapy;
2. Conventional therapy.

When clinical stability is achieved, the patient is randomized in either the NT-proBNP-titrated group or the conventional therapy group. In the NT-proBNP-titrated group a NT-proBNP reduction of > 30% before discharge is pursued. When patients do not reach the >30% NT-proBNP reduction, they follow a predefined algorithm before discharge to improve the number of patients discharged with >30% reduction in NT-proBNP levels. Hence, duration of admission can be prolonged with a maximum of 2-4 days. The NT-proBNP levels of patients in the conventional treatment group will be measured but not revealed to patients, physicians or nurses. Conventional treatment encompasses the recommended therapy depicted in the current guidelines for acute heartfailure.

## **Contactpersonen**

### **Publiek**

Academic Medical Center "C <br>  
University of Amsterdam<br>  
Department of cardiology.  
Wouter E.M. Kok  
Amsterdam  
The Netherlands  
+31 (0)20 5665777

### **Wetenschappelijk**

Academic Medical Center "C <br>  
University of Amsterdam<br>  
Department of cardiology.  
Wouter E.M. Kok  
Amsterdam  
The Netherlands  
+31 (0)20 5665777

# Deelname eisen

## Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Hospital admission because of clinically validated acutely decompensated heart failure. A clinical diagnosis of HF is made on the basis of a HF-score of 2 points or higher;
2. Elevated NT-proBNP levels  $\geq 1700$  ng/L ( $\geq 200$  pmol/L) on hospital admission;
3. Written informed consent to participate in this study prior to any study procedures.

## Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Severe Chronic Obstructive Pulmonary Disease (COPD) with  $FEV_1 < 1$  l/min;
2. Pulmonary embolism within 1 month prior to admission and pulmonary hypertension not caused by left ventricle dysfunction (LVD);
3. Patients undergoing Continuous Ambulant Peritoneal Dialysis (CAPD)/ Haemodialysis patients;
4. Patients with planned Coronary Artery Bypass Grafting (CABG), Percutaneous Coronary Intervention (PCI), Cardiac Resynchronization Therapy (CRT) and/or valvular surgery before admission (until one day before admission);
5. Patients with planned Coronary Artery Bypass Grafting (CABG), Percutaneous Coronary Intervention (PCI), Cardiac Resynchronization Therapy (CRT) and/or valvular surgery during admission until the moment of randomization;
6. Patient with a history of ST-segment-Elevated Myocardial Infarction (STEMI), CABG, PCI, CRT and/or valvular surgery within 1 month prior to admission;
7. Signed informed consent for any current interventional study;
8. Presence of severe non-cardiac related life-threatening disease before inclusion with an expected survival of less than 6 months after inclusion;
9. Mental or physical status not allowing written informed consent;
10. Unwillingness to give informed consent;

11. Circumstances that prevent follow-up (no permanent home address, transient, etc.).

## Onderzoeksopzet

### Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

### Deelname

Nederland	
Status:	Werving gestopt
(Verwachte) startdatum:	01-11-2011
Aantal proefpersonen:	400
Type:	Werkelijke startdatum

## Ethische beoordeling

Positief advies	
Datum:	08-02-2012
Soort:	Eerste indiening

## Registraties

### Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 44072  
Bron: ToetsingOnline  
Titel:

## Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

## In overige registers

Register	ID
NTR-new	NL3128
NTR-old	NTR3279
CCMO	NL36873.018.11
ISRCTN	ISRCTN wordt niet meer aangevraagd.
OMON	NL-OMON44072

## Resultaten

### Samenvatting resultaten

N/A