

Effects of preventive case management on parenting and children of mental patients.

Gepubliceerd: 12-10-2010 Laatste bijgewerkt: 18-08-2022

Parents with psychiatric problems often have parenting problems, in particular when they have to deal with other risk factors for poor parenting like for instance isolation and poverty. This means an enhanced risk for their children on developing...

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON22367

Bron

Nationaal Trial Register

Aandoening

psychiatric disorders, depression, prevention, COPMI

Ondersteuning

Primaire sponsor: ZonMW; Fonds Nuts Ohra; Parnassia Bavo Group

Overige ondersteuning: ZonMW; Fonds Nuts Ohra; Parnassia Bavo Group

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Parenting, measured by the HOME Inventory (, retrieved 26-10-2010) and by the Parenting Daily Hassles (Cronic & Greenberg, 1990).

Toelichting onderzoek

Achtergrond van het onderzoek

Effects of preventive casemanagement on parenting of mental patients and behavioral problems in children is studied in a RCT with 116 families. Consumption of preventive and indicated care is used for a CEA.

Doel van het onderzoek

Parents with psychiatric problems often have parenting problems, in particular when they have to deal with other risk factors for poor parenting like for instance isolation and poverty. This means an enhanced risk for their children on developing behavioral problems. Coordinated early help for these high risk families as offered by the program Preventive Basic Care Management (PBCM), could prevent severe parenting problems and thereby reduce the risk of behavioural problems in the children.

Onderzoeksopzet

Baseline, 9 months, 18 months:

1. HOME, Parenting Daily Hassles;
2. SDQ;
3. Questionnaire Support and Help.

Onderzoeksproduct en/of interventie

The PBCM intervention includes:

1. Systematic assessment;
2. Intervention planning and coordinating supportive services, tailored to address the identified risk factors of poor parenting;
3. Monitoring and evaluating the implementation of the indicated supportive services and their effects.

Assessment:

The assessment is standardised and focused on (1) identifying early signs of behavioural problems in children, (2) description and evaluation of parenting behaviour, and (3) taxation

of risk and protective factors for parenting, including parental competences. Risk assessment is done in home visits, interviews with parents and involved services, such as schools, child care centres, youth health services and therapists. The assessment is used to identify needs for preventive support in parents and children. Parents are explicitly asked about the goals they want to achieve through their involvement in the PBCM programme. These goals are central targets for the next step.

Intervention planning and coordination:

This assessment is used to design a tailored intervention plan for the family. The plan is developed in several steps. First, the assessment is discussed in the PBCM team to design a draft intervention plan. Then, the assessment and this intervention proposal are discussed with (preferably both) parents and involved services in a joint meeting.

First priorities in the PBCM process often address parenting problems with structuring daily family life and parental dissatisfaction with psychiatric treatment or stagnations. Families are referred to Family Services for help with improving daily structure and organisation of family life. The PBCM manager discusses how treatment stagnations can be solved with the therapist and the parent. Secondly, interventions for tackling contextual risk factors and improving contextual protective factors such as co-parent arrangement, social support, family living conditions, utilization of Child Care, poverty are planned.

The final plan is family tailored, assessment-based and consists of multiple interventions. Actions are formulated in terms of parenting behaviours and actions of services. The PBCM manager sets criteria for the goal and the implementation of services in order to adjust them to the needs and risks of the family. Timing and dosage are optimally adjusted to the capacities of the parents. He/she advises parents, facilitates the use of specialised services and advocates their interest. Finally, he/she gives an overview of the agreed actions, and documents concrete settlements. Clear goals and settlements are written down and sent to all participants.

Monitoring and evaluation:

Follow up co-ordination meetings with parents and services are planned bi-monthly and constitute a core element in the Case Management strategy. In these bi-monthly meetings the PBCM manager discusses the progress and the settlements made in the previous meeting in a systematic way.

The programme ends when a sufficient level of positive parenting behaviour is accomplished and the targeted reduction of risk factors is secured. The maximum duration of the programme is eighteen months. As a rule families have about eight meetings during this period.

The control condition includes a flyer about effects of psychiatric problems of parents in children and the possibility for personal advice and supportgroups for children and/or parents of the COPMI-programme. COPMI stands for Children of Parents with a Mental Illness. Parents are free to participate in these and other preventive services on their own initiative, but therapists will not actively or systematically stimulate the use of services.

Contactpersonen

Publiek

Lijnbaan 4
Henny Wansink
Parnassia Bavo Group, Prevention Department
Den Haag 2512 VA
The Netherlands
+31 (0)70 3918389

Wetenschappelijk

Lijnbaan 4
Henny Wansink
Parnassia Bavo Group, Prevention Department
Den Haag 2512 VA
The Netherlands
+31 (0)70 3918389

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Psychiatric patients with children between 3 and 9 years (not suffering from a childhood mental disorder or mental retardation);
2. Parenting problems;
3. More than three risk factors which threaten good parenting.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. No psychiatric treatment or finish of treatment within three months;
2. Therapist is referring child to social services.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-09-2010
Aantal proefpersonen:	116
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	12-10-2010
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL2453
NTR-old	NTR2569
Ander register	ZonMw : 80-82435-98-9125
ISRCTN	ISRCTN wordt niet meer aangevraagd.

Resultaten

Samenvatting resultaten

N/A