

The effects of protein type and added leucine on post-exercise muscle protein synthesis

Gepubliceerd: 12-03-2015 Laatste bijgewerkt: 15-05-2024

Protein-carbohydrate ingestion stimulates greater muscle protein synthesis than carbohydrate placebo alone. milk and whey protein elicit the greatest anabolic response. Soy protein with added leucine will stimulate muscle protein synthesis...

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON23128

Bron

Nationaal Trial Register

Verkorte titel

PepsiCo

Aandoening

Young healthy men

Ondersteuning

Primaire sponsor: NUTRIM School for Nutrition, Toxicology, and Metabolism Maastricht University

Overige ondersteuning: PepsiCo

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Toelichting onderzoek

Achtergrond van het onderzoek

Rationale: Dietary protein intake after exercise is necessary to maximally stimulate muscle protein synthesis rates. Data demonstrate that the type/source of protein consumed (e.g., animal vs. plant-derived proteins) can impact the amplitude and duration of muscle protein synthesis during post-exercise recovery. Specifically, bovine milk proteins stimulate greater rates of muscle protein synthesis after resistance exercise than consumption of an isonitrogenous soy-protein beverage. The major proteins in bovine milk are casein (~80%) and whey protein (~20%). In their isolated forms, these proteins greatly differ in their digestion and absorption kinetics. Whey protein is rapidly digested and absorbed leading to a pronounced, rapid, but transient peak in plasma amino acid levels and robust stimulation of protein synthesis. On the other hand, isolated casein is a slowly digested protein that results in a slower, moderate, but more prolonged increase in plasma amino acid availability resulting in a greatly attenuated protein synthesis response compared to whey. No studies have compared the muscle protein synthesis response following bovine milk as compared to its constituent proteins whey and casein. Further, whether the post-exercise muscle protein synthesis response to soy protein can be enhanced when the leucine content of soy is matched to milk remains unknown. We aim to fill this gap in our understanding.

Objective: To define the properties of whey, casein, milk protein, as well as soy protein with and without additional leucine to augment post-exercise muscle protein synthesis when co-ingested with a carbohydrate containing recovery drink.

Study design: Parallel group, randomized, placebo controlled, double blind.

Study population: 72 young (20-30 y inclusive) healthy males.

Intervention: Subjects will perform both aerobic and resistance exercise and consume either a carbohydrate solution or a carbohydrate solution with 20 g whey, 20 g casein, 20 g milk protein, 20 g soy protein, or 20 g soy protein with leucine. In addition, continuous intravenous tracer infusions will be applied, with plasma and muscle samples collected.

Main study parameters/endpoints: Primary: mixed, myofibrillar, and mitochondrial protein bound [13C6] phenylalanine enrichments. Secondary: plasma glucose, insulin, leucine, phenylalanine, tyrosine, plasma [13C6]phenylalanine and (3,5-D2)-tyrosine enrichments.

Doel van het onderzoek

Protein-carbohydrate ingestion stimulates greater muscle protein synthesis than carbohydrate placebo alone. milk and whey protein elicit the greatest anabolic response. Soy protein with added leucine will stimulate muscle protein synthesis equivalent to whey protein

Onderzoeksopzet

t=0 Drink. t=0 , t=2h and t=6h muscle biopsies. 14 blood draws

Onderzoeksproduct en/of interventie

Exercise bout followed by ingestion of one of the following beverages:

- Carbohydrate drink (45 g Carbohydrate)
- Carbohydrate drink (45 g Carbohydrate) with 20 g milk protein
- Carbohydrate drink (45 g Carbohydrate) with 20 g whey
- Carbohydrate drink (45 g Carbohydrate) with 20 g casein
- Carbohydrate drink (45 g Carbohydrate) with 20 g soy
- Carbohydrate drink (45 g Carbohydrate) with 20 g soy and leucine

Contactpersonen

Publiek

Department of Human Movement Sciences - Maastricht University Medical Centre+

T.A. Churchward-Venne
P.O. Box 616

Maastricht 6200 MD
The Netherlands
+31 43 388 2154

Wetenschappelijk

Department of Human Movement Sciences - Maastricht University Medical Centre+

T.A. Churchward-Venne
P.O. Box 616

Maastricht 6200 MD
The Netherlands
+31 43 388 2154

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

- Males
- Aged between 20-30 years
- Bodyweight between 65-95 kg inclusive
- Healthy, recreationally active
- BMI < 25 kg/m²

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

- The use of over-the-counter nutritional supplements excluding multivitamins/minerals
- Having any identified metabolic or intestinal disorders
- Use of tobacco products
- Non-steroidal anti-inflammatory drugs (NSAID) in the 4 days prior to the experimental trial
- Allergies to milk proteins (whey or casein) or soy protein
- Vegetarians
- Arthritic conditions
- A history of neuromuscular problems
- Previous participation in amino acid tracer studies
- Individuals on any medications known to affect protein metabolism (i.e. corticosteroids, non-steroidal anti-inflammatories, or prescription strength acne medications).
- Diabetes

- Training more than 4 days per week

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Parallel
Toewijzing:	Gerandomiseerd
Blinding:	Dubbelblind
Controle:	Actieve controle groep

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-09-2014
Aantal proefpersonen:	72
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	12-03-2015
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 42133
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL4844
NTR-old	NTR5098
CCMO	NL49732.068.14
OMON	NL-OMON42133

Resultaten