

Outpatient group SFT for BPD.

Gepubliceerd: 02-12-2007 Laatste bijgewerkt: 15-05-2024

Group SFT treatment is more effective than individual SFT treatment for Borderline Personality Disorder in an outpatient setting.

Ethische beoordeling	Niet van toepassing
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON26741

Bron

Nationaal Trial Register

Verkorte titel

GROST-BOR study

Aandoening

1. Borderline personality disorder;
2. group therapy;
3. schema focused therapy;
4. treatment effectivity.

(NLD: borderline persoonlijkheidsstoornis, groepsbehandeling, schemagerichte therapie, behandel-effectiviteit).

Ondersteuning

Primaire sponsor: Research school experimental psychopathology university Maastricht.

Overige ondersteuning: Research school experimental psychopathology university Maastricht, ministry of OC&W.

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

Severity of Borderline Personality symptoms (BPDSI-IV-, semi-structured interview) measuring the current severity and frequency of the DSM-IV BPD manifestations.

Toelichting onderzoek

Achtergrond van het onderzoek

The present study is a continuation of a previous study on the effectivity of outpatient treatment of Borderline Personality Disorder (BPD) by means of Schema Focused Therapy (SFT) as presented by Giesen-Bloo (2006). In this study SFT proved to be effective in reducing borderline personality disorder-specific and general psychopathologic dysfunctioning and in improving quality of life. Due to long duration of the therapy, treatment is costly and waitinglists are long. Because of these reasons SFT has been implemented in group therapy in several institutions for mental health care in the Netherlands. As far as we know, the effectivity of this group-SFT has not been studied so far. In this study group SFT therapy will be compared with individual SFT therapy.

Doel van het onderzoek

Group SFT treatment is more effective than individual SFT treatment for Borderline Personality Disorder in an outpatient setting.

Onderzoeksopzet

Primary outcome measurement at baseline, every six months, until six months post-treatment (6 measurements).

Onderzoeksproduct en/of interventie

individual SFT (historical data) versus group schemafocused therapy (with a duration of two years) for patients with borderline personality disorder in an outpatient setting.

Contactpersonen

Publiek

parallelweg 45-47

V.E.M Dickhaut
Maastricht 6221 BD
The Netherlands
043-3299640

Wetenschappelijk

parallelweg 45-47

V.E.M Dickhaut
Maastricht 6221 BD
The Netherlands
043-3299640

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

1. Main diagnosis of borderline personality disorder (BPDSI-IV score greater than 20);
2. 18-60 years of age;
3. Dutch language.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

1. Psychotic disorder;
2. subthreshold anti social personality disorder or narcissistic personality disorder;
3. bipolar disorder;
4. dissociative identity disorder;
5. attention deficit/hyperactivity disorder;

6. addiction that requires detoxification;

7. mental retardation.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Factorieel
Toewijzing:	Niet-gerandomiseerd
Blinding:	Open / niet geblindeerd
Controle:	N.v.t. / onbekend

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	01-03-2008
Aantal proefpersonen:	20
Type:	Verwachte startdatum

Ethische beoordeling

Niet van toepassing	
Soort:	Niet van toepassing

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

ID: 31733
Bron: ToetsingOnline
Titel:

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1121
NTR-old	NTR1156
CCMO	NL20841.068.07
ISRCTN	ISRCTN wordt niet meer aangevraagd
OMON	NL-OMON31733

Resultaten

Samenvatting resultaten

Giesen-Bloo, J. et al. (2006), Outpatient psychotherapy for borderline personality disorder: Randomized trial of schema-focused therapy vs transference focused psychotherapy. Archives of General Psychiatry, 63, 649-658.