

The effects of high-protein breakfasts.

Gepubliceerd: 16-10-2007 Laatste bijgewerkt: 18-08-2022

The administration of a whey protein/ alphalactalbumine/ green tea mixture to a standardized yogurt breakfast, will cause an increase in post-prandial energy expenditure, due to diet-induced thermogenesis and an increase in satiety within 5 hours.

Ethische beoordeling	Positief advies
Status	Werving gestart
Type aandoening	-
Onderzoekstype	Interventie onderzoek

Samenvatting

ID

NL-OMON28372

Bron

Nationaal Trial Register

Verkorte titel

The effects of high protein breakfasts.

Aandoening

1. Voedselingredienten;
2. energiegebruik;
3. verzadiging;
4. positieve energiebalans;
5. food ingredients;
6. energy expenditure;
7. satiety;
8. positive energy balance.

Ondersteuning

Primaire sponsor: Friesland Nutrition

PO Box 226

8901 MA Leeuwarden

Phone: +31582992434

Fax: +31582992286

E-mail: lucie.vanderzee@frieslandfoods.com

Overige ondersteuning: Friesland Nutrition

Onderzoeksproduct en/of interventie

Uitkomstmaten

Primaire uitkomstmaten

1. Energy expenditure;
2. Satiety, using the Visual Analogue Scale (VAS) score (0 to 10, 0= no hunger and 10= extreme hunger).

Toelichting onderzoek

Achtergrond van het onderzoek

High protein diets cause a thermogenic effect and increase satiety. When people are satiated, energy intake will be restricted. A positive energy balance, can be prevented because of the addition of extra proteins to the breakfasts. During the next meal energy intake will be less, because people might be satiated and still full. Also the energy expenditure will be increased by the extra proteins which also prevents a positive energy balance.

The additional green tea during one of the interventions also causes a thermogenic effect. Green tea contains different components which inhibit particular enzymes. Through this the sympathetic activity is increased and also the energy expenditure.

Doel van het onderzoek

The administration of a whey protein/ algalactalbumine/ green tea mixture to a standardized yogurt breakfast, will cause an increase in post-prandial energy expenditure, due to diet-induced thermogenesis and an increase in satiety within 5 hours.

Onderzoeksopzet

One testday is 5 hours. 30 minutes before the intervention the energy expenditure is

measured and after the intervention energy expenditure is measured for 4,5 hours. Satiety is measured every hour and 1 hour after the intervention even every 20 minutes.

Onderzoeksproduct en/of interventie

Subjects will eat 4 different breakfasts. Subjects will come to the university once a week for 4 consecutive weeks. Their energy expenditure will be measured by means of indirect calorimetry (ventilated hood). Also, satiety is measured with visual analog scales. Randomization will determine in which sequence the subjects will receive the interventions. These are the interventions:

1. placebo yogurt (15% proteins, 47% carbohydrates and 38% fat);
2. placebo yogurt + whey protein (41% proteins, 47% carbohydrates and 12% fat);
3. placebo yogurt + algalactalbumine (41% proteins, 47% carbohydrates and 12% fat);
4. placebo yogurt + green tea catechins (15% proteins, 47% carbohydrates and 38% fat).

The breakfast consists of 20% of the daily energy intake and is subject specific.

Contactpersonen

Publiek

Maastricht University
Dept. Humane Biology
PO Box 616

Rick Hursel
Maastricht 6200 MD
The Netherlands
+31 43 3882123

Wetenschappelijk

Maastricht University
Dept. Humane Biology
PO Box 616

Rick Hursel

Deelname eisen

Belangrijkste voorwaarden om deel te mogen nemen (Inclusiecriteria)

Healthy men and women, with a Body Mass Index (BMI) between 22-33 kg/m², and age of 20-60 years.

Belangrijkste redenen om niet deel te kunnen nemen (Exclusiecriteria)

Exclusion criteria for subjects are apart from age and BMI:

1. smoking;
2. having food allergies (especially milk protein allergies);
3. being on medication (except the use of contraception);
4. excessive alcohol consumption;
5. excessive exercise;
6. not being weight stable;
7. being dietary restraint (assessed by the Three Eating Questionnaire (TFEQ));
8. people who drink more than one cup of coffee per day.

Onderzoeksopzet

Opzet

Type:	Interventie onderzoek
Onderzoeksmodel:	Cross-over
Toewijzing:	Gerandomiseerd
Blinding:	Enkelblind
Controle:	Placebo

Deelname

Nederland	
Status:	Werving gestart
(Verwachte) startdatum:	07-01-2008
Aantal proefpersonen:	40
Type:	Verwachte startdatum

Ethische beoordeling

Positief advies	
Datum:	16-10-2007
Soort:	Eerste indiening

Registraties

Opgevolgd door onderstaande (mogelijk meer actuele) registratie

Geen registraties gevonden.

Andere (mogelijk minder actuele) registraties in dit register

Geen registraties gevonden.

In overige registers

Register	ID
NTR-new	NL1062
NTR-old	NTR1095
Ander register	MEC nr. : 07-3-080.

Register

ISRCTN

ID

ISRCTN wordt niet meer aangevraagd

Resultaten

Samenvatting resultaten

N/A